

LAMBERT SERIES IN ANALYTIC NUMBER THEORY

JORDAN BELL

ABSTRACT. Tour of 19th and early 20th century analytic number theory.

1. INTRODUCTION

Let $d(n)$ denote the number of positive divisors of n . For $|z| < 1$,

$$\sum_{n=1}^{\infty} d(n)z^n = \sum_{n=1}^{\infty} \frac{z^n}{1-z^n}.$$

2. EULER

The first use of the term “Lambert series” was by Euler to describe the roots of an equation.

Euler writes in E25 [28] about the particular value of a Lambert series.

3. LAMBERT

Lambert [53, pp. 506–511, §875]

Monatsbuch, September 1764, “Singula haec in Capp. ult. Ontol. occurrunt”,
and Anm. 5, Anm. 25, 1764, Anm. 12 1765, Anm. 19, 1765 [2].

Youschkevitch [87]

Bullynck [7]

Lorey [59, p. 23]

Löwenhaupt [60, p. 32]

4. KRAFFT

Krafft [50, pp. 244–245]

5. SERVOIS

Servois [72] and [73, p. 166].

6. LACROIX

Lacroix [51, pp. 465–466, §1195]

7. KLÜGEL

Klügel [46, pp. 52–53, s.v. “Theiler einer Zahl”, §12]:

Ist $N = \alpha^m \beta^n \gamma^p \dots$, wo α, β, γ , Primzahlen sind; so erhellet auch leicht, daß alle Theiler von N , die Einheit und die Zahl selbst mit eingeschlossen, durch die Glieder des Products

$$(1 + \alpha + \alpha^2 + \dots + \alpha^m)(1 + \beta + \beta^2 + \dots + \beta^n)(1 + \gamma + \gamma^2 + \dots + \gamma^p) \dots$$

argestellt werden. Die Anzahl der Glieder dieses Products, d. i. die Anzahl aller Theiler von N , ist offenbar $= (m + 1)(n + 1)(p + 1) \dots$. Für das obige Beispiel $= 4 \cdot 3 \cdot 2 = 24$, wo die Einheit mit eingeschlossen ist.

In der aus der Entwicklung von

$$\frac{x}{1-x} + \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} + \dots + \frac{x^n}{1-x^n} + \dots$$

entspringenden Reihe:

$$x + 2x^2 + 2x^3 + 3x^4 + 2x^5 + 4x^6 + 2x^7 + \dots$$

welche Lambert in seiner Architektonik S. 507. mittheilt, enthält jeder Coefficient so viele Einheiten, als der Exponent der entsprechenden Potenz von x Theiler hat.

8. STERN

Stern [77]

9. CLAUSEN

Clausen [21] states that

$$\sum_{n=1}^{\infty} \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} x^{n^2} \left(\frac{1+x^n}{1-x^n} \right),$$

and that the right-hand series converges quickly for small x . Clausen does prove this expansion, and a proof is later given by Scherk [68]. Scherk’s argument uses the fact

$$1 + 2t + 2t^2 + 2t^3 + 2t^4 + \dots = (1 + t + t^2 + t^3 + t^4 + \dots) + t(1 + t + t^2 + t^3 + t^4 + \dots) = \frac{1+t}{1-t}.$$

We write

$$\sum_{n=1}^{\infty} \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} x^{nm}.$$

The series is

x	$+x^2$	$+x^3$	$+x^4$	$+x^5$	$+x^6$	$+etc.$
$+x^2$	$+x^4$	$+x^6$	$+x^8$	$+x^{10}$	$+x^{12}$	$+etc.$
$+x^3$	$+x^6$	$+x^9$	$+x^{12}$	$+x^{15}$	$+x^{18}$	$+etc.$
$+x^4$	$+x^8$	$+x^{12}$	$+x^{16}$	$+x^{20}$	$+x^{24}$	$+etc.$
$+x^5$	$+x^{10}$	$+x^{15}$	$+x^{20}$	$+x^{25}$	$+x^{30}$	$+etc.$
$+x^6$	$+x^{12}$	$+x^{18}$	$+x^{24}$	$+x^{30}$	$+x^{36}$	$+etc.$
$+etc.$						

We sum the terms in the first row and column: the sum of these is

$$x + 2x^2 + 2x^3 + 2x^4 + \text{etc.} = x \left(\frac{1+x}{1-x} \right).$$

Then, from what remains we sum the terms in the second row and column: the sum of these is

$$x^4 + 2x^6 + 2x^8 + 2x^{10} + \text{etc.} = x^4 \left(\frac{1+x^2}{1-x^2} \right).$$

Then, from what remains, we sum the terms in the third row and column: the sum of these is

$$x^9 + 2x^{12} + 2x^{15} + 2x^{18} + \text{etc.} = x^9 \left(\frac{1+x^3}{1-x^3} \right),$$

etc.

10. EISENSTEIN

Eisenstein [27] states that for $|z| < 1$,

$$\sum_{n=1}^{\infty} \frac{z^n}{1-z^n} = \frac{1}{(1-x)(1-x^2)(1-x^3)\dots} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{nz^{n(n+1)/2}}{(1-x)\dots(1-x^n)}.$$

For $t = \frac{1}{z}$, Eisenstein states that

$$\frac{z}{1-z} + \frac{z^2}{1-z^2} + \frac{z^3}{1-z^3} + \frac{z^4}{1-z^4} + \text{etc.}$$

is equal to

$$\frac{1}{t-1 - \frac{(t-1)^2}{t^2-1 - \frac{t(t-1)^2}{t^3-1 - \frac{t^2(t^2-1)^2}{t^4-1 - \frac{t^2(t^3-1)^2}{t^5-1 - \frac{t^3(t^3-1)^2}{t^6-1 - \frac{t^7-1}{\text{etc.}}}}}}}}$$

Expressing Lambert series using continued fractions is relevant to the irrationality of the value of the series. See Borwein [3]. See also Zudilin [90].

11. MÖBIUS

Möbius [62]

12. JACOBI

Jacobi's *Fundamenta nova* [44, §40, 66 and p. 185]

Chandrasekharan [20, Chapter X]: using Lambert series to prove the four squares theorem.

13. DIRICHLET

Dirichlet [25]

Fischer [29]

14. CAUCHY

Cauchy [13] and [14] two memoirs in the same volume.

15. BURHENNE

Burhenne [8] says the following about Lambert series. For

$$F(x) = \sum_{n=1}^{\infty} d(n)x^n,$$

we have

$$d(n) = \frac{F^{(n)}(0)}{n!}.$$

Define

$$F_k(x) = \frac{x^k}{1-x^k},$$

so that

$$F(x) = \sum_{k=1}^{\infty} F_k(x).$$

It is apparent that if $k > n$, then

$$F_k^{(n)}(0) = 0,$$

hence

$$F^{(n)}(0) = \sum_{k=1}^n F_k^{(n)}(0).$$

The above suggests finding explicit expressions for $F_k^{(n)}(0)$. Burhenne cites Sohncke [74, pp. 32–33]: for even k ,

$$\begin{aligned} \frac{d^n \left(\frac{x^p}{x^k - a^k} \right)}{dx^n} &= (-1)^n \frac{n!}{ka^{k-p-1}} \left(\frac{1}{(x-a)^{n+1}} - (-1)^p \frac{1}{(x+a)^{n+1}} \right) \\ &+ (-1)^n \frac{n!}{\frac{1}{2}ka^{k-p-1}} \sum_{h=1}^{\frac{1}{2}k-1} \frac{\cos \left(\frac{2h(p+1)\pi}{k} + (n+1)\phi_h \right)}{\sqrt{(x^2 - 2xa \cos \frac{2h\pi}{n} + a^2)^{n+1}}} \end{aligned}$$

and for odd k ,

$$\begin{aligned} \frac{d^n \left(\frac{x^p}{x^k - a^k} \right)}{dx^n} &= (-1)^n \frac{n!}{ka^{k-p-1}} \frac{1}{(x-a)^{n+1}} \\ &+ (-1)^n \frac{n!}{\frac{1}{2}ka^{k-p-1}} \sum_{h=1}^{\frac{k-1}{2}} \frac{\cos \left(\frac{2h(p+1)\pi}{k} + (n+1)\phi_h \right)}{\sqrt{(x^2 - 2xa \cos \frac{2h\pi}{n} + a^2)^{n+1}}}, \end{aligned}$$

where

$$\cos \phi_h = \frac{x - a \cos \frac{2h\pi}{k}}{\sqrt{x^2 - 2xa \cos \frac{2h\pi}{k} + a^2}}, \quad \sin \phi_h = \frac{a \sin \frac{2h\pi}{k}}{\sqrt{x^2 - 2xa \cos \frac{2h\pi}{k} + a^2}}.$$

For $a = 1$ and $x = 0$,

$$\cos \phi_h = -\cos \frac{2h\pi}{k}, \quad \sin \phi_h = \sin \frac{2h\pi}{k},$$

from which

$$\phi_h = \pi - \frac{2h\pi}{k},$$

and thus

$$\begin{aligned} \cos\left(\frac{2h(k+1)\pi}{k} + (n+1)\phi_h\right) &= \cos\left(\frac{2h(k+1)\pi}{k} + (n+1)\left(\pi - \frac{2h\pi}{k}\right)\right) \\ &= \cos\left(2h\pi + \frac{2h\pi}{k} + \pi - \frac{2h\pi}{k} + n\left(\pi - \frac{2h\pi}{k}\right)\right) \\ &= \cos\left((n+1)\pi - \frac{2nh\pi}{k}\right) \\ &= (-1)^{n+1} \cos\frac{2nh\pi}{k}. \end{aligned}$$

For even k , taking $p = k$ we have

$$\frac{d^n\left(\frac{x^k}{1-x^k}\right)}{dx^n} = (-1)^{n+1} \frac{n!}{k} \left(\frac{1}{(-1)^{n+1}} - 1\right) + (-1)^{n+1} \frac{n!}{\frac{1}{2}k} \sum_{h=1}^{\frac{1}{2}k-1} (-1)^{n+1} \cos\frac{2nh\pi}{k},$$

i.e.,

$$F_k^{(n)}(0) = \frac{n!}{k} (1 - (-1)^{n+1}) + \frac{2 \cdot n!}{k} \sum_{h=1}^{\frac{1}{2}k-1} \cos\frac{2nh\pi}{k}.$$

For odd k , taking $p = k$ we have

$$\frac{d^n\left(\frac{x^k}{1-x^k}\right)}{dx^n} = (-1)^{n+1} \frac{n!}{k} \frac{1}{(-1)^{n+1}} + (-1)^{n+1} \frac{n!}{\frac{1}{2}k} \sum_{h=1}^{\frac{k-1}{2}} (-1)^{n+1} \cos\frac{2nh\pi}{k},$$

i.e.,

$$F_k^{(n)}(0) = \frac{n!}{k} + \frac{2 \cdot n!}{k} \sum_{h=1}^{\frac{k-1}{2}} \cos\frac{2nh\pi}{k}.$$

Using the identity, for $h \notin 2\pi\mathbb{Z}$,

$$\sum_{h=1}^M \cos h\theta = -\frac{1}{2} + \frac{\sin\left(M + \frac{1}{2}\right)\theta}{2\sin\frac{\theta}{2}} = -\frac{1}{2} + \frac{1}{2} \left(\sin M\theta \cot\frac{\theta}{2} + \cos M\theta\right),$$

we get for even k ,

$$\begin{aligned} F_k^{(n)}(0) &= \begin{cases} \frac{n!}{k} \cot\frac{n\pi}{k} \sin n\pi & k \nmid n \\ \frac{n!}{k} (1 - (-1)^{n+1}) + \frac{2 \cdot n!}{k} \left(\frac{1}{2}k - 1\right) & k|n \end{cases} \\ &= \begin{cases} 0 & k \nmid n \\ n! - \frac{n!}{k} (1 + (-1)^{n+1}) & k|n. \end{cases} \end{aligned}$$

For odd k ,

$$\begin{aligned} F_k^{(n)}(0) &= \begin{cases} \frac{n!}{k} \csc\frac{n\pi}{k} \sin n\pi & k \nmid n \\ \frac{n!}{k} + \frac{2 \cdot n!}{k} \frac{k-1}{2} & k|n. \end{cases} \\ &= \begin{cases} 0 & k \nmid n \\ n! & k|n. \end{cases} \end{aligned}$$

16. ZEHFUSS

Zehfuss [88]

17. BERNOULLI NUMBERS

The **Bernoulli polynomials** are defined by

$$\frac{te^{tx}}{e^t - 1} = \sum_{m=0}^{\infty} B_m(x) \frac{t^m}{m!}.$$

The **Bernoulli numbers** are defined by $B_m = B_m(0)$.

We denote by $[x]$ the greatest integer $\leq x$, and we define $\{x\} = x - [x]$, namely, the fractional part of x . We define $P_m(x) = B_m(\{x\})$, the **periodic Bernoulli functions**.

18. EULER-MACLAURIN SUMMATION FORMULA

Euler E47 and E212, §142, for the summation formula. Euler's studies the gamma function in E368. In particular, in §12 he gives Stirling's formula, and in §14 he obtains $\Gamma'(1) = -\gamma$. Euler in §142 of E212 states that

$$\gamma = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{(-1)^{n+1} B_{2n}}{2n}.$$

Bromwich [6, Chapter XII]

The Euler-Maclaurin summation formula [5, p. 280, Ch. VI, Eq. 35] tells us that for $f \in C^\infty([0, 1])$,

$$f(0) = \int_0^1 f(t) dt + B_1(f(1) - f(0)) + \sum_{m=1}^k \frac{1}{(2m)!} B_{2m}(f^{(2m-1)}(1) - f^{(2m-1)}(0)) + R_{2k},$$

where

$$R_{2k} = - \int_0^1 \frac{P_{2k}(1-\eta)}{(2k)!} f^{(2k)}(\eta) d\eta.$$

Poisson and Jacobi on the Euler-Maclaurin summation formula.

19. SCHLÖMILCH

Schlömilch [69] and [71, p. 238], [70]

For $m \geq 1$,

$$(1) \quad \int_0^\infty \frac{t^{2m-1}}{e^{2\pi t} - 1} dt = (-1)^{m+1} \frac{B_{2m}}{4m}.$$

For $\alpha > 0$,

$$(2) \quad \int_0^\infty \frac{\sin \alpha t}{e^{2\pi t} - 1} dt = \frac{1}{4} + \frac{1}{2} \left(\frac{1}{e^\alpha - 1} - \frac{1}{\alpha} \right)$$

and

$$(3) \quad \int_0^\infty \frac{1 - \cos \alpha t}{e^{2\pi t} - 1} \frac{dt}{t} = \frac{1}{4} \alpha + \frac{1}{2} (\log(1 - e^{-\alpha}) - \log \alpha).$$

For $\xi > 0$ and $n \geq 1$, using (2) with $\alpha = \xi, 2\xi, 3\xi, \dots, 2n\xi$ and also using

$$\sum_{k=1}^N \sin k\theta = \frac{1}{2} \cot \frac{\theta}{2} - \frac{\cos(N + \frac{1}{2})\theta}{2 \sin \frac{\theta}{2}},$$

we get

$$\begin{aligned} \sum_{m=1}^{2n} \left(\frac{1}{e^{m\xi} - 1} - \frac{1}{m\xi} \right) &= \sum_{m=1}^{2n} \left(-\frac{1}{2} + 2 \int_0^\infty \frac{\sin m\xi t}{e^{2\pi t} - 1} dt \right) \\ &= -n + \int_0^\infty \frac{1}{e^{2\pi t} - 1} \sum_{m=1}^{2n} 2 \sin m\xi t dt \\ &= -n + \int_0^\infty \frac{1}{e^{2\pi t} - 1} \left(\cot \frac{\xi t}{2} - \frac{\cos(2n + \frac{1}{2})\xi t}{\sin \frac{\xi t}{2}} \right) dt. \end{aligned}$$

Using $\cos(a + b) = \cos a \cos b - \sin a \sin b$, this becomes

$$(4) \quad \sum_{m=1}^{2n} \left(\frac{1}{e^{m\xi} - 1} - \frac{1}{m\xi} \right) = -n + \int_0^\infty \frac{1}{e^{2\pi t} - 1} (1 - \cos 2n\xi t) \cot \frac{\xi t}{2} dt \\ + \int_0^\infty \frac{1}{e^{2\pi t} - 1} \sin 2n\xi t dt.$$

For $\alpha = 2n\xi$, (3) tells us

$$\int_0^\infty \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} \frac{dt}{t} = \frac{1}{4} \cdot 2n\xi + \frac{1}{2} (\log(1 - e^{-2n\xi}) - \log 2n\xi).$$

Rearranging,

$$(5) \quad \frac{\log 2n}{\xi} = n + \frac{\log(1 - e^{-2n\xi}) - \log \xi}{\xi} - \frac{2}{\xi} \int_0^\infty \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} \frac{dt}{t}$$

Adding (4) and (5) gives

$$\begin{aligned} &\sum_{m=1}^{2n} \frac{1}{e^{m\xi} - 1} - \frac{1}{\xi} \left(-\log 2n + \sum_{m=1}^{2n} \frac{1}{m} \right) \\ &= \frac{\log(1 - e^{-2n\xi}) - \log \xi}{\xi} - \int_0^\infty \left(\frac{2}{\xi t} - \cot \frac{\xi t}{2} \right) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ &\quad + \int_0^\infty \frac{1}{e^{2\pi t} - 1} \sin 2n\xi t dt. \end{aligned}$$

Writing

$$C_n = -\log n + \sum_{m=1}^n \frac{1}{m}$$

and using (2) this becomes

$$\begin{aligned} & \sum_{m=1}^{2n} \frac{1}{e^{m\xi} - 1} - \frac{1}{\xi} C_{2n} \\ &= \frac{\log(1 - e^{-2n\xi}) - \log \xi}{\xi} - 2 \int_0^\infty \left(\frac{1}{\xi t} - \frac{1}{2} \cot \frac{\xi t}{2} \right) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ & \quad + \frac{1}{4} + \frac{1}{2} \left(\frac{1}{e^{2n\xi} - 1} - \frac{1}{2n\xi} \right). \end{aligned}$$

We write

$$I_{2n}(\xi) = 2 \int_0^\infty \left(\frac{1}{\xi t} - \frac{1}{2} \cot \frac{\xi t}{2} \right) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt,$$

and we shall obtain an asymptotic formula for $I_{2n}(\xi)$.

We apply the Euler-Maclaurin summation formula. Let $h > 0$, and for $f(t) = \cos ht$ we have $f'(t) = -h \sin ht$, and for $m \geq 1$ we have $f^{(2m)}(t) = (-1)^m h^{2m} \cos ht$ and $f^{(2m-1)}(t) = (-1)^m h^{2m-1} \sin ht$. Thus the Euler-Maclaurin formula yields

$$1 = \int_0^1 \cos ht dt - \frac{1}{2}(\cos h - 1) + \sum_{m=1}^k \frac{1}{(2m)!} B_{2m} (-1)^m h^{2m-1} \sin h + R_{2k}.$$

Using the identity $\cot \frac{\theta}{2} = \frac{1 + \cos \theta}{\sin \theta}$ and dividing by $\sin h$, this becomes

$$(6) \quad \frac{1}{2} \cot \frac{h}{2} = \frac{1}{h} + \sum_{m=1}^k \frac{1}{(2m)!} B_{2m} (-1)^m h^{2m-1} + \frac{1}{\sin h} R_{2k}.$$

Because $P_m(1 - \eta) = P_m(\eta)$ for even m ,

$$\begin{aligned} R_{2k} &= - \int_0^1 \frac{P_{2k}(\eta)}{(2k)!} (-1)^k h^{2k} \cos h\eta d\eta \\ &= -B_{2k} \int_0^1 \frac{1}{(2k)!} (-1)^k h^{2k} \cos h\eta d\eta - \int_0^1 \frac{(P_{2k}(\eta) - B_{2k})}{(2k)!} (-1)^k h^{2k} \cos h\eta d\eta \\ &= (-1)^{k+1} \frac{B_{2k} h^{2k}}{(2k)!} \frac{\sin h}{h} + (-1)^{k+1} \frac{h^{2k}}{(2k)!} \int_0^1 (P_{2k}(\eta) - B_{2k}) \cos h\eta d\eta. \end{aligned}$$

Since $P_{2k}(\eta) - B_{2k}$ does not change sign on $(0, 1)$, by the mean-value theorem for integration there is some $\theta = \theta(h, k)$, $0 < \theta < 1$, such that (using $\int_0^1 P_{2k}(\eta) d\eta = 0$)

$$\int_0^1 (P_{2k}(\eta) - B_{2k}) \cos h\eta d\eta = \cos h\theta \int_0^1 (P_{2k}(\eta) - B_{2k}) d\eta = -B_{2k} \cos h\theta.$$

Therefore (6) becomes

$$\begin{aligned} \frac{1}{2} \cot \frac{h}{2} - \frac{1}{h} &= \sum_{m=1}^k \frac{1}{(2m)!} B_{2m} (-1)^m h^{2m-1} \\ & \quad + (-1)^{k+1} \frac{B_{2k} h^{2k-1}}{(2k)!} + (-1)^{k+2} \frac{h^{2k}}{(2k)! \sin h} B_{2k} \cos h\theta, \end{aligned}$$

i.e.,

$$\frac{1}{2} \cot \frac{h}{2} - \frac{1}{h} = \sum_{m=1}^{k-1} \frac{1}{(2m)!} B_{2m} (-1)^m h^{2m-1} + (-1)^k \frac{h^{2k}}{(2k)! \sin h} B_{2k} \cos h\theta.$$

Write

$$E_k(h) = (-1)^{k+1} \frac{h^{2k}}{(2k)! \sin h} B_{2k} \cos h\theta.$$

We apply the above to $I_{2n}(\xi)$, and get, for any $k \geq 1$,

$$\begin{aligned} I_{2n}(\xi) &= 2 \int_0^\infty \left(E_k(\xi t) - \sum_{m=1}^{k-1} \frac{1}{(2m)!} B_{2m} (-1)^m (\xi t)^{2m-1} \right) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ &= -2 \sum_{m=1}^{k-1} \frac{1}{(2m)!} B_{2m} (-1)^m \xi^{2m-1} \int_0^\infty t^{2m-1} \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ &\quad + 2 \int_0^\infty E_k(\xi t) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt. \end{aligned}$$

Using (1),

$$\begin{aligned} \int_0^\infty t^{2m-1} \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt &= \int_0^\infty \frac{t^{2m-1}}{e^{2\pi t} - 1} dt - \int_0^\infty \frac{t^{2m-1} \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ &= (-1)^{m+1} \frac{B_{2m}}{4m} - \int_0^\infty \frac{t^{2m-1} \cos 2n\xi t}{e^{2\pi t} - 1} dt. \end{aligned}$$

Let

$$f(x) = \frac{1}{e^x - 1} - \frac{1}{x}.$$

By (2),

$$f(x) + \frac{1}{2} = 2 \int_0^\infty \frac{\sin xt}{e^{2\pi t} - 1} dt.$$

For $m \geq 1$,

$$f^{(2m-1)}(x) = 2 \int_0^\infty \frac{(-1)^{m-1} t^{2m-1} \cos xt}{e^{2\pi t} - 1} dt,$$

which for $x = 2n\xi$ becomes

$$\frac{(-1)^{m-1}}{2} f^{(2m-1)}(2n\xi) = \int_0^\infty \frac{t^{2m-1} \cos 2n\xi t}{e^{2\pi t} - 1} dt.$$

Therefore

$$2 \int_0^\infty t^{2m-1} \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt = (-1)^{m+1} \frac{B_{2m}}{2m} + (-1)^m f^{(2m-1)}(2n\xi).$$

Thus $I_{2n}(\xi)$ is

$$\begin{aligned} I_{2n}(\xi) &= - \sum_{m=1}^{k-1} \frac{1}{(2m)!} B_{2m} (-1)^m \xi^{2m-1} \left((-1)^{m+1} \frac{B_{2m}}{2m} + (-1)^m f^{(2m-1)}(2n\xi) \right) \\ &\quad + 2 \int_0^\infty E_k(\xi t) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ &= \sum_{m=1}^{k-1} \frac{B_{2m}^2}{(2m)! 2m} \xi^{2m-1} - \sum_{m=1}^{k-1} \frac{B_{2m}}{(2m)!} \xi^{2m-1} f^{(2m-1)}(2n\xi) \\ &\quad + 2 \int_0^\infty E_k(\xi t) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt. \end{aligned}$$

But

$$\begin{aligned} \left| \int_0^\infty E_k(\xi t) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \right| &= \left| \int_0^\infty (-1)^{k+1} \frac{(\xi t)^{2k}}{(2k)! \sin \xi t} B_{2k} \cos \xi t \theta \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \right| \\ &\leq \frac{|B_{2k}|}{(2k)!} \int_0^\infty \frac{(\xi t)^{2k}}{|\sin \xi t|} \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt. \end{aligned}$$

It is a fact that for all $u \in \mathbb{R}$,

$$\frac{1 - \cos 2nu}{|\sin u|} \leq \frac{\pi}{2} \frac{1 - \cos 2nu}{u},$$

we obtain

$$\begin{aligned} &\left| \int_0^\infty E_k(\xi t) \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \right| \\ &\leq \frac{\pi |B_{2k}|}{2 (2k)!} \int_0^\infty (\xi t)^{2k-1} \frac{1 - \cos 2n\xi t}{e^{2\pi t} - 1} dt \\ &= \frac{\pi |B_{2k}|}{2 (2k)!} \xi^{2k-1} \cdot \frac{1}{2} \left((-1)^{k+1} \frac{B_{2k}}{2k} + (-1)^k f^{(2k-1)}(2n\xi) \right). \end{aligned}$$

Hence

$$\begin{aligned} I_{2n}(\xi) &= \sum_{m=1}^{k-1} \frac{B_{2m}^2}{(2m)! 2m} \xi^{2m-1} - \sum_{m=1}^{k-1} \frac{B_{2m}}{(2m)!} \xi^{2m-1} f^{(2m-1)}(2n\xi) \\ &\quad + O\left(\frac{B_{2k}^2}{(2k)! 2k} \xi^{2k-1}\right) + O\left(\frac{|B_{2k}|}{(2k)!} \xi^{2k-1} f^{(2k-1)}(2n\xi)\right). \end{aligned}$$

Therefore we have

$$\begin{aligned} &\sum_{m=1}^{2n} \frac{1}{e^{m\xi} - 1} - \frac{1}{\xi} C_{2n} \\ &= \frac{\log(1 - e^{-2n\xi}) - \log \xi}{\xi} + \frac{1}{4} + \frac{1}{2} \left(\frac{1}{e^{2n\xi} - 1} - \frac{1}{2n\xi} \right) - I_{2n}(\xi) \\ &= \frac{\log(1 - e^{-2n\xi}) - \log \xi}{\xi} + \frac{1}{4} + \frac{1}{2} \left(\frac{1}{e^{2n\xi} - 1} - \frac{1}{2n\xi} \right) \\ &\quad - \sum_{m=1}^{k-1} \frac{B_{2m}^2}{(2m)! 2m} \xi^{2m-1} + \sum_{m=1}^{k-1} \frac{B_{2m}}{(2m)!} \xi^{2m-1} f^{(2m-1)}(2n\xi) \\ &\quad + O\left(\frac{B_{2k}^2}{(2k)! 2k} \xi^{2k-1}\right) + O\left(\frac{|B_{2k}|}{(2k)!} \xi^{2k-1} f^{(2k-1)}(2n\xi)\right). \end{aligned}$$

Taking $n \rightarrow \infty$,

$$\sum_{m=1}^{\infty} \frac{1}{e^{m\xi} - 1} - \frac{\gamma}{\xi} = -\frac{\log \xi}{\xi} + \frac{1}{4} - \sum_{m=1}^{k-1} \frac{B_{2m}^2}{(2m)! 2m} \xi^{2m-1} + O\left(\frac{B_{2k}^2}{(2k)! 2k} \xi^{2k-1}\right).$$

20. VORONOI SUMMATION FORMULA

The Voronoi summation formula [22, p. 182] states that if $f : \mathbb{R} \rightarrow \mathbb{C}$ is a Schwartz function, then

$$\begin{aligned} \sum_{n=1}^{\infty} d(n)f(n) &= \int_0^{\infty} f(t)(\log t + 2\gamma)dt + \frac{f(0)}{4} \\ &\quad + \sum_{n=1}^{\infty} d(n) \int_0^{\infty} f(t)(4K_0(4\pi(nt)^{1/2}) - 2\pi Y_0(4\pi(nt)^{1/2}))dt, \end{aligned}$$

where K_0 and Y_0 are Bessel functions.

Let $0 < x < 1$. For $f(t) = e^{-tx}$, we compute

$$\begin{aligned} &\int_0^{\infty} f(t)(4K_0(4\pi(nt)^{1/2}) - 2\pi Y_0(4\pi(nt)^{1/2}))dt \\ &= -\frac{2}{x} \exp\left(\frac{4\pi^2 n}{x}\right) \text{Ei}\left(-\frac{4\pi^2 n}{x}\right) - \frac{2}{x} \exp\left(-\frac{4\pi^2 n}{x}\right) \text{Ei}\left(\frac{4\pi^2 n}{x}\right), \end{aligned}$$

where

$$\text{Ei}(x) = -\int_{-x}^{\infty} \frac{e^{-t}}{t} dt, \quad x \neq 0,$$

the exponential integral. Then the Voronoi summation formula yields

$$\begin{aligned} &\sum_{n=1}^{\infty} d(n)e^{-nx} \\ &= \frac{\gamma}{x} - \frac{\log x}{x} + \frac{1}{4} \\ &\quad + \sum_{n=1}^{\infty} d(n) \left(-\frac{2}{x} \exp\left(\frac{4\pi^2 n}{x}\right) \text{Ei}\left(-\frac{4\pi^2 n}{x}\right) - \frac{2}{x} \exp\left(-\frac{4\pi^2 n}{x}\right) \text{Ei}\left(\frac{4\pi^2 n}{x}\right) \right). \end{aligned}$$

Egger and Steiner [26] give a proof of the Voronoi summation formula involving Lambert series.

Kluyver [47] and [48]

Guinand [36]

21. CURTZE

Curtze [23]

22. LAGUERRE

Laguerre [52]

23. V. A. LEBESGUE

V. A. Lebesgue [56]:

24. BOUNIAKOWSKY

Bouniakowsky [4]

25. CHEBYSHEV

Chebyshev [80]

26. CATALAN

Catalan [9]
Catalan [10, p. 89]
Catalan [11, p. 119, §CXXIV] and [12, pp. 38–39, §CCXXVI]

27. PINCHERLE

Pincherle [63]

28. GLAISHER

Glaisher [34, p. 163]

29. GÜNTHER

Günther [37, p. 83] and [38, p. 178]

30. STIELTJES

Stieltjes [78]
cf. Zhang [89]

31. ROGEL

Rogel [65] and [66]

32. CESÀRO

Cesàro [15]
Cesàro [16]
Cesàro [17] and [18, pp. 181–184]
Bromwich [6, p. 201, Chapter VIII, Example B, 35]

33. DE LA VALLÉE-POUSSIN

de la Vallée-Poussin [24]

34. TORELLI

Torelli [83]

35. FIBONACCI NUMBERS

Landau [54]

36. KNOPP

Knopp [49]

37. GENERATING FUNCTIONS

Hardy and Wright [41, p. 258, Theorem 307]:

Theorem 1. For $f(s) = \sum_{n=1}^{\infty} a_n n^{-s}$ and $g(s) = \sum_{n=1}^{\infty} b_n n^{-s}$,

$$\sum_{n=1}^{\infty} a_n \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} b_n x^n, \quad |x| < 1,$$

if and only if there is some σ such that

$$\zeta(s)f(s) = g(s), \quad \operatorname{Re}(s) > \sigma.$$

For $f(s) = \sum_{n=1}^{\infty} \mu(n)n^{-s}$ and $g(s) = 1$, using [41, p. 250, Theorem 287]

$$\frac{1}{\zeta(s)} = \sum_{n=1}^{\infty} \mu(n)n^{-s}, \quad \operatorname{Re}(s) > 1,$$

we get

$$(7) \quad \sum_{n=1}^{\infty} \frac{\mu(n)x^n}{1-x^n} = x.$$

For $f(s) = \sum_{n=1}^{\infty} \phi(n)n^{-s}$ and

$$g(s) = \zeta(s-1) = \sum_{n=1}^{\infty} n^{-s+1} = \sum_{n=1}^{\infty} nn^{-s},$$

using [41, p. 250, Theorem 288]

$$\frac{\zeta(s-1)}{\zeta(s)} = \sum_{n=1}^{\infty} \phi(n)n^{-s}, \quad \operatorname{Re}(s) > 2,$$

we get

$$\sum_{n=1}^{\infty} \frac{\phi(n)x^n}{1-x^n} = \sum_{n=1}^{\infty} nx^n = \frac{x}{(1-x)^2}.$$

For $n = p_1^{a_1} \cdots p_r^{a_r}$, define $\Omega(n) = a_1 + \cdots + a_n$ and

$$\lambda(n) = (-1)^{\Omega(n)}.$$

For $f(s) = \sum_{n=1}^{\infty} \lambda(n)n^{-s}$ and

$$g(s) = \zeta(2s) = \sum_{n=1}^{\infty} n^{-2s} = \sum_{n=1}^{\infty} (n^2)^{-s},$$

using [41, p. 255, Theorem 300]

$$\frac{\zeta(2s)}{\zeta(s)} = \sum_{n=1}^{\infty} \lambda(n)n^{-s}, \quad \operatorname{Re}(s) > 1,$$

we get

$$\sum_{n=1}^{\infty} \frac{\lambda(n)x^n}{1-x^n} = \sum_{n=1}^{\infty} x^{n^2}.$$

We define the **von Mangoldt function** $\Lambda : \mathbb{N} \rightarrow \mathbb{R}$ by $\Lambda(n) = \log p$ if n is some positive integer power of a prime p , and $\Lambda(n) = 0$ otherwise. For example,

$\Lambda(1) = 0$, $\Lambda(12) = 0$, $\Lambda(125) = \log 5$. It is a fact [41, p. 254, Theorem 296] that for any n , the von Mangoldt function satisfies

$$(8) \quad \sum_{m|n} \Lambda(m) = \log n.$$

For $f(s) = \sum_{n=1}^{\infty} \Lambda(n)n^{-s}$ and

$$g(s) = -\zeta'(s) = \sum_{n=1}^{\infty} \log nn^{-s},$$

using [41, p. 253, Theorem 294]

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \Lambda(n)n^{-s},$$

we obtain

$$\sum_{n=1}^{\infty} \frac{\Lambda(n)x^n}{1-x^n} = \sum_{n=1}^{\infty} \log nx^n.$$

38. MERTENS

For $\operatorname{Re} s > 1$, we define

$$P(s) = \sum_p \frac{1}{p^s}.$$

We also define

$$H = \sum_{m=2}^{\infty} \sum_p \frac{1}{mp^m}.$$

Mertens [61] proves the following.

Theorem 2. *As $\varrho \rightarrow 0$,*

$$P(1 + \varrho) = \log \left(\frac{1}{\varrho} \right) - H + o(1).$$

Proof. As $\varrho \rightarrow 0$,

$$\zeta(1 + \varrho) = \frac{1}{\varrho} + \gamma + O(\varrho) = \frac{1}{\varrho}(1 + \gamma\varrho + O(\varrho^2)).$$

Taking the logarithm,

$$(9) \quad \log \zeta(1 + \varrho) = \log \left(\frac{1}{\varrho} \right) + \log(1 + \gamma\varrho + O(\varrho^2)) = \log \left(\frac{1}{\varrho} \right) + \gamma\varrho + O(\varrho^2).$$

On the other hand, for $\varrho > 0$,

$$\zeta(1 + \varrho) = \prod_p \frac{1}{1 - \frac{1}{p^{1+\varrho}}},$$

and taking the logarithm,

$$\begin{aligned} \log \zeta(1 + \varrho) &= - \sum_p \log \left(1 - \frac{1}{p^{1+\varrho}} \right) \\ &= \sum_p \sum_{m=1}^{\infty} \frac{1}{mp^{m(1+\varrho)}} \\ &= P(1 + \rho) + \sum_{m=2}^{\infty} \sum_p \frac{1}{mp^{m(1+\varrho)}}. \end{aligned}$$

Then as $\varrho \rightarrow 0$,

$$\log \zeta(1 + \varrho) = P(1 + \varrho) + H + o(1).$$

Combining this with (9) we get that as $\varrho \rightarrow 0$,

$$P(1 + \rho) = \log \left(\frac{1}{\rho} \right) - H + o(1).$$

□

Mertens [61] also proves that for any x there is some

$$|\delta| < \frac{4}{\log(x+1)} + \frac{2}{x \log x}$$

such that

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + \gamma - H + \delta.$$

Thus,

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + \gamma - H + O\left(\frac{1}{\log x}\right).$$

Mertens shows that

$$H = - \sum_{n=2}^{\infty} \mu(n) \frac{\log \zeta(n)}{n}.$$

This can be derived using (7), and we do this now; see [58].

Lemma 3. For $\operatorname{Re} s > 1$,

$$\frac{1}{s} \log \zeta(s) = \int_2^{\infty} \frac{\pi(t) dt}{t(t^s - 1)}.$$

Proof. For p prime and $\operatorname{Re} s > 0$,

$$\begin{aligned}
\int_p^\infty \frac{dt}{t(t^s - 1)} &= \int_p^\infty t^{-s-1} \frac{1}{1 - t^{-s}} dt \\
&= \int_p^\infty t^{-s-1} \sum_{n=0}^{\infty} (t^{-s})^n dt \\
&= \sum_{n=0}^{\infty} \int_p^\infty t^{-ns-s-1} dt \\
&= \sum_{n=0}^{\infty} \left. \frac{t^{-ns-s}}{-ns-s} \right|_p^\infty \\
&= \frac{1}{s} \sum_{n=1}^{\infty} \frac{p^{-ns}}{n} \\
&= -\frac{1}{s} \log(1 - p^{-s}),
\end{aligned}$$

hence

$$\log\left(\frac{1}{1 - p^{-s}}\right) = s \int_p^\infty \frac{dt}{t(t^s - 1)}.$$

On the one hand,

$$\sum_p \int_p^\infty \frac{dt}{t(t^s - 1)} = \int_2^\infty \frac{\pi(t) dt}{t(t^s - 1)}.$$

On the other hand, for $\operatorname{Re} s > 1$ we have

$$\sum_p \log\left(\frac{1}{1 - p^{-s}}\right) = \log \prod_p \left(\frac{1}{1 - p^{-s}}\right) = \log \zeta(s).$$

Combining these, for $\operatorname{Re} s > 1$,

$$\frac{1}{s} \log \zeta(s) = \int_2^\infty \frac{\pi(t) dt}{t(t^s - 1)}.$$

□

Theorem 4.

$$H = -\sum_{n=2}^{\infty} \mu(n) \frac{\log \zeta(n)}{n}.$$

Proof. For any prime p and for $m \geq 1$,

$$\int_p^\infty t^{-m-1} dt = \left. \frac{t^{-m}}{-m} \right|_p^\infty = \frac{1}{mp^m},$$

and using this we have

$$\begin{aligned}
H &= \sum_{m=2}^{\infty} \sum_p \frac{1}{mp^m} \\
&= \sum_{m=2}^{\infty} \sum_p \int_p^{\infty} t^{-m-1} dt \\
&= \sum_{m=2}^{\infty} \int_2^{\infty} \pi(t) t^{-m-1} dt \\
&= \int_2^{\infty} \pi(t) \left(\sum_{m=2}^{\infty} t^{-m-1} \right) dt \\
&= \int_2^{\infty} \pi(t) \frac{1}{t^2(t-1)} dt
\end{aligned}$$

Rearranging (7),

$$\frac{x^2}{1-x} = - \sum_{n=2}^{\infty} \frac{\mu(n)x^n}{1-x^n}.$$

With $x = t^{-1}$,

$$\frac{1}{t(t-1)} = - \sum_{n=2}^{\infty} \frac{\mu(n)}{t^n - 1},$$

so

$$\frac{1}{t^2(t-1)} = - \sum_{n=2}^{\infty} \frac{\mu(n)}{t(t^n - 1)}.$$

Thus we have

$$H = - \int_2^{\infty} \pi(t) \left(\sum_{n=2}^{\infty} \frac{\mu(n)}{t(t^n - 1)} \right) dt = - \sum_{n=2}^{\infty} \mu(n) \int_2^{\infty} \frac{\pi(t) dt}{t(t^n - 1)}.$$

Using Lemma 3 for $s = 2, 3, 4, \dots$,

$$H = - \sum_{n=2}^{\infty} \mu(n) \cdot \frac{1}{n} \log \zeta(n),$$

completing the proof. \square

39. PRELIMINARIES ON PRIME NUMBERS

We define

$$\vartheta(x) = \sum_{p \leq x} \log p = \log \prod_{p \leq x} p$$

and

$$\psi(x) = \sum_{p^m \leq x} \log p = \sum_{n \leq x} \Lambda(n).$$

One sees that

$$\psi(x) = \sum_{p \leq x} [\log_p x] \log p = \sum_{p \leq x} \left[\frac{\log x}{\log p} \right] \log p.$$

As well,

$$(10) \quad \psi(x) = \sum_{m=1}^{\infty} \sum_{p \leq x^{1/m}} \log p = \sum_{m=1}^{\infty} \vartheta(x^{1/m});$$

there are only finitely many terms on the right-hand side, as $\vartheta(x^{1/m}) = 0$ if $x < 2^m$.

Theorem 5.

$$\psi(x) = \vartheta(x) + O(x^{1/2}(\log x)^2).$$

Proof. For $x \geq 2$, $\vartheta(x) < x \log x$, giving

$$\begin{aligned} \sum_{2 \leq m \leq \frac{\log x}{\log 2}} \vartheta(x^{1/m}) &< \sum_{2 \leq m \leq \frac{\log x}{\log 2}} x^{1/m} \frac{1}{m} \log x \\ &\leq x^{1/2} \log x \sum_{2 \leq m \leq \frac{\log x}{\log 2}} \frac{1}{m} \\ &= O(x^{1/2}(\log x)^2). \end{aligned}$$

Thus, using (10) we have

$$\psi(x) = \vartheta(x) + \sum_{2 \leq m \leq \frac{\log x}{\log 2}} \vartheta(x^{1/m}) = \vartheta(x) + O(x^{1/2}(\log x)^2).$$

□

We prove that if $\lim_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = 1$ then $\frac{\pi(x)}{x/\log x} = 1$.

Theorem 6.

$$\liminf_{x \rightarrow \infty} \frac{\pi(x)}{x/\log x} = \liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x}$$

and

$$\limsup_{x \rightarrow \infty} \frac{\pi(x)}{x/\log x} = \limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x}.$$

Proof. From (10), $\vartheta(x) \leq \psi(x)$. And,

$$\psi(x) = \sum_{p \leq x} \left[\frac{\log x}{\log p} \right] \log p \leq \sum_{p \leq x} \frac{\log x}{\log p} \log p = \log x \sum_{p \leq x}.$$

Hence

$$\frac{\vartheta(x)}{x} \leq \frac{\pi(x) \log x}{x},$$

whence

$$\liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \leq \liminf_{x \rightarrow \infty} \frac{\pi(x)}{x/\log x}$$

and

$$\limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \leq \limsup_{x \rightarrow \infty} \frac{\pi(x)}{x/\log x}.$$

Let $0 < \alpha < 1$. For $x > 1$,

$$\vartheta(x) = \sum_{p \leq x} \log p \geq \sum_{x^\alpha < p \leq x} \log p > \sum_{x^\alpha < p \leq x} \log x^\alpha = \alpha \log x (\pi(x) - \pi(x^\alpha)).$$

As $\pi(x^\alpha) < x^\alpha$,

$$\vartheta(x) > \alpha\pi(x) \log x - \alpha x^\alpha \log x,$$

i.e.,

$$\frac{\vartheta(x)}{x} > \alpha \frac{\pi(x) \log x}{x} - \alpha \frac{\log x}{x^{1-\alpha}}.$$

This yields

$$\liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \geq \alpha \liminf_{x \rightarrow \infty} \frac{\pi(x) \log x}{x} - \alpha \liminf_{x \rightarrow \infty} \frac{\log x}{x^{1-\alpha}} = \alpha \liminf_{x \rightarrow \infty} \frac{\pi(x) \log x}{x}$$

and

$$\limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \geq \alpha \limsup_{x \rightarrow \infty} \frac{\pi(x) \log x}{x} - \alpha \limsup_{x \rightarrow \infty} \frac{\log x}{x^{1-\alpha}} = \alpha \limsup_{x \rightarrow \infty} \frac{\pi(x) \log x}{x}.$$

Since these are true for all $0 < \alpha < 1$, we obtain respectively

$$\liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \geq \liminf_{x \rightarrow \infty} \frac{\pi(x) \log x}{x}$$

and

$$\limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \geq \limsup_{x \rightarrow \infty} \frac{\pi(x) \log x}{x}.$$

□

40. WIENER'S TAUBERIAN THEOREM

Wiener [85, Chapter III].

Wiener-Ikehara [19]

Rudin [67, p. 229, Theorem 9.7]

We say that a function $s : (0, \infty) \rightarrow \mathbb{R}$ is **slowly decreasing** if

$$\liminf(s(\rho v) - s(v)) \geq 0, \quad v \rightarrow \infty, \quad \rho \rightarrow 1^+.$$

Widder [84, p. 211, Theorem 10b]: Wiener's tauberian theorem tells us that if $a \in L^\infty(0, \infty)$ and is slowly decreasing and if $g \in L^1(0, \infty)$ satisfies

$$\int_0^\infty t^{ix} g(t) dt \neq 0, \quad x \in \mathbb{R},$$

then

$$\lim_{x \rightarrow \infty} \frac{1}{x} \int_0^\infty g\left(\frac{t}{x}\right) a(t) dt = A \int_0^\infty g(t) dt$$

implies that

$$\lim_{v \rightarrow \infty} a(v) = A.$$

It is straightforward to check the following by rearranging summation.

Lemma 7. *If $\sum_{n=1}^\infty a_n z^n$ has radius of convergence ≥ 1 , then for $|z| < 1$,*

$$\sum_{n=1}^\infty a_n \frac{z^n}{1-z^n} = \sum_{n=1}^\infty \left(\sum_{m|n} a_m \right) z^n.$$

Using Lemma 7 with $a_n = \Lambda(n)$ and $z = e^{-x}$ and applying (8), we get

$$(11) \quad \sum_{n=1}^{\infty} \Lambda(n) \frac{z^n}{1-z^n} = \sum_{n=1}^{\infty} \log(n) z^n.$$

From (11), and Lemma 7 with $a_n = 1$, we have

$$\sum_{n=1}^{\infty} (\Lambda(n) - 1) \frac{e^{-nx}}{1-e^{-nx}} = \sum_{n=1}^{\infty} (\log n - d(n)) e^{-nx}.$$

We follow Widder [84, p. 231, Theorem 16.6].

Theorem 8. *As $x \rightarrow 0^+$,*

$$\sum_{n=1}^{\infty} (\log n - d(n)) e^{-nx} = -\frac{2\gamma}{x} + O(x^{-1/2}).$$

Proof. Generally,

$$\begin{aligned} (1-z) \sum_{n=1}^{\infty} z^n \sum_{m=1}^n a_m &= (1-z) \sum_{m=1}^{\infty} a_m \sum_{n=m}^{\infty} z^n \\ &= (1-z) \sum_{m=1}^{\infty} a_m \frac{z^m}{1-z} \\ &= \sum_{m=1}^{\infty} a_m z^m. \end{aligned}$$

Using this with $a_m = \log m - d(m)$ and $z = e^{-x}$ gives

$$\begin{aligned} \sum_{n=1}^{\infty} (\log n - d(n)) e^{-nx} &= (1-e^{-x}) \sum_{n=1}^{\infty} e^{-nx} \left(\sum_{m=1}^n \log m - \sum_{m=1}^n d(m) \right) \\ &= (1-e^{-x}) \sum_{n=1}^{\infty} e^{-nx} \left(\log(n!) - \sum_{m=1}^n d(m) \right). \end{aligned}$$

Using

$$\log(n!) = n \log n - n + O(\log n)$$

and

$$\sum_{m=1}^n d(m) = n \log n + (2\gamma - 1)n + O(n^{1/2}),$$

we get

$$\log(n!) - \sum_{m=1}^n d(m) = -2\gamma n + O(n^{1/2}).$$

Therefore,

$$\sum_{n=1}^{\infty} (\log n - d(n)) e^{-nx} = (1-e^{-x}) \sum_{n=1}^{\infty} e^{-nx} (-2\gamma n + O(n^{1/2})).$$

One proves that there is some K such that for all $0 \leq y < 1$,

$$(1-y) \left(\log \frac{1}{y} \right)^{1/2} \sum_{n=1}^{\infty} n^{1/2} y^n \leq K,$$

whence, with $y = e^{-x}$,

$$\sum_{n=1}^{\infty} n^{1/2} e^{-nx} \leq K \frac{x^{-1/2}}{1 - e^{-x}}.$$

Also,

$$\sum_{n=1}^{\infty} n e^{-nx} = \frac{e^{-x}}{(1 - e^{-x})^2},$$

and thus we have

$$\begin{aligned} \sum_{n=1}^{\infty} (\log n - d(n)) e^{-nx} &= -2\gamma \frac{e^{-x}}{1 - e^{-x}} + O(x^{-1/2}) \\ &= -2\gamma \frac{1}{e^x - 1} + O(x^{-1/2}). \end{aligned}$$

But

$$\frac{1}{e^x - 1} = \frac{1}{x} - \frac{1}{2} + O(x),$$

so

$$\sum_{n=1}^{\infty} (\log n - d(n)) e^{-nx} = -\frac{2\gamma}{x} + O(x^{-1/2}).$$

□

Define

$$f(x) = \sum_{n=1}^{\infty} (\Lambda(n) - 1) \frac{e^{-nx}}{1 - e^{-nx}},$$

and

$$h(x) = \sum_{n \leq x} \frac{\Lambda(n) - 1}{n},$$

and

$$g(t) = \frac{d}{dt} \left(\frac{te^{-t}}{1 - e^{-t}} \right).$$

First we show that h is slowly decreasing.

Lemma 9. $h(x)$ is slowly decreasing.

Proof. Using

$$\sum_{1 \leq n \leq x} \frac{1}{n} = \log x + \gamma + O(x^{-1}), \quad x \rightarrow \infty,$$

we have, for $0 < x < \infty$ and $\rho > 1$,

$$\begin{aligned} h(\rho x) - h(x) &= \sum_{x < n \leq \rho x} \frac{\Lambda(n) - 1}{n} \\ &\geq - \sum_{x < n \leq \rho x} \frac{1}{n} \\ &= - \sum_{1 \leq n \leq \rho x} \frac{1}{n} + \sum_{1 \leq n \leq x} \frac{1}{n} \\ &= -\log(\rho x) + \log x + O((\rho x)^{-1}) + O(x^{-1}) \\ &= -\log \rho + O((\rho x)^{-1}) + O(x^{-1}). \end{aligned}$$

Hence as $x \rightarrow \infty$ and $\rho \rightarrow 1^+$,

$$h(\rho x) - h(x) \rightarrow 0,$$

which shows that h is slowly decreasing. □

The following is from Widder [84, pp. 231–232].

Lemma 10. *As $x \rightarrow \infty$,*

$$\frac{1}{x} \int_0^\infty g\left(\frac{t}{x}\right) h(t) dt = 2\gamma + O(x^{-1/2}).$$

Proof. Let $I(t) = 0$ for $t < 0$ and $I(t) = 1$ for $t \geq 0$. Writing

$$h(x) = \sum_{n=1}^{\infty} I(x-n) \frac{\Lambda(n) - 1}{n},$$

we check that for $x > 0$,

$$\begin{aligned} \int_0^\infty \frac{te^{-xt}}{1-e^{-xt}} dh(t) &= \sum_{n=1}^{\infty} \int_0^\infty \frac{te^{-xt}}{1-e^{-xt}} \frac{\Lambda(n) - 1}{n} d(I(t-n)) \\ &= \sum_{n=1}^{\infty} \int_0^\infty \frac{te^{-xt}}{1-e^{-xt}} \frac{\Lambda(n) - 1}{n} d\delta_n(t) \\ &= \sum_{n=1}^{\infty} \frac{ne^{-nx}}{1-e^{-nx}} \frac{\Lambda(n) - 1}{n} \\ &= f(x). \end{aligned}$$

On the other hand, integrating by parts,

$$\begin{aligned} f(x) &= \int_0^\infty \frac{te^{-xt}}{1-e^{-xt}} dh(t) \\ &= \int_0^\infty \frac{1}{x} \frac{xt e^{-xt}}{1-e^{-xt}} dh(t) \\ &= \int_0^\infty \frac{1}{x} \frac{xt e^{-xt}}{1-e^{-xt}} dh(t) \\ &= \int_0^\infty \frac{1}{x} \frac{te^{-t}}{1-e^{-t}} dh\left(\frac{t}{x}\right) \\ &= \frac{1}{x} \frac{te^{-t}}{1-e^{-t}} h\left(\frac{t}{x}\right) \Big|_0^\infty - \int_0^\infty \frac{1}{x} g(t) h\left(\frac{t}{x}\right) dt \\ &= - \int_0^\infty \frac{1}{x} g(t) h\left(\frac{t}{x}\right) dt \\ &= - \int_0^\infty g(xt) h(t) dt. \end{aligned}$$

By Theorem 8, as $x \rightarrow 0^+$,

$$f(x) = -\frac{2\gamma}{x} + O(x^{-1/2}),$$

i.e., as $x \rightarrow 0^+$,

$$\int_0^\infty g(xt) h(t) dt = \frac{2\gamma}{x} + O(x^{-1/2}).$$

Thus, as $x \rightarrow \infty$,

$$\int_0^\infty g\left(\frac{t}{x}\right) h(t) dt = 2\gamma x + O(x^{1/2}).$$

□

The following is from Widder [84, p. 232].

Lemma 11.

$$\int_0^\infty t^{-ix} g(t) dt = \begin{cases} -1 & x = 0 \\ ix\zeta(1-ix)\Gamma(1-ix) & x \neq 0. \end{cases}$$

Proof.

$$\begin{aligned} \int_0^\infty t^{-ix} g(t) dt &= \int_0^\infty t^{-ix} \frac{d}{dt} \left(\frac{te^{-t}}{1-e^{-t}} \right) dt \\ &= \lim_{\delta \rightarrow 0} \int_0^\infty t^{-ix+\delta} \frac{d}{dt} \left(\frac{te^{-t}}{1-e^{-t}} \right) dt \\ &= \lim_{\delta \rightarrow 0} \left(t^{-ix+\delta} \frac{te^{-t}}{1-e^{-t}} \Big|_0^\infty + (ix-\delta) \int_0^\infty t^{-ix+\delta-1} \frac{te^{-t}}{1-e^{-t}} dt \right) \\ &= \lim_{\delta \rightarrow 0} (ix-\delta) \int_0^\infty t^{-ix+\delta-1} \frac{te^{-t}}{1-e^{-t}} dt \\ &= \lim_{\delta \rightarrow 0} (ix-\delta) \int_0^\infty \frac{t^{(-ix+\delta+1)-1} e^{-t}}{1-e^{-t}} dt. \end{aligned}$$

Using

$$\int_0^\infty \frac{t^{s-1}}{e^t-1} dt = \zeta(s)\Gamma(s), \quad \operatorname{Re}(s) > 1,$$

this becomes

$$\int_0^\infty t^{-ix} g(t) dt = \lim_{\delta \rightarrow 0^+} (ix-\delta)\zeta(1+\delta-ix)\Gamma(1+\delta-ix).$$

If $x = 0$, then using

$$\zeta(s) = \frac{1}{s-1} + \gamma + O(|s-1|), \quad s \rightarrow 1,$$

we get

$$\lim_{\delta \rightarrow 0^+} (-\delta)\zeta(1+\delta)\Gamma(1+\delta) = -1.$$

If $x > 0$, then

$$\lim_{\delta \rightarrow 0^+} (ix-\delta)\zeta(1+\delta-ix)\Gamma(1+\delta-ix) = ix\zeta(1-ix)\Gamma(1-ix).$$

□

By Wiener's tauberian theorem, it follows that

$$\sum_{n=1}^\infty \frac{\Lambda(n) - 1}{n} = -2\gamma.$$

Lemma 12.

$$h(x) = \int_{\frac{1}{2}}^x \frac{d(\psi(t) - [t])}{t}.$$

Proof. Let $I(t) = 0$ for $t < 0$ and $I(t) = 1$ for $t \geq 0$. Writing

$$\psi(x) = \sum_{n=1}^{\infty} I(x-n)\Lambda(n), \quad [x] = \sum_{n=1}^{\infty} I(x-n),$$

we have

$$\begin{aligned} \int_{\frac{1}{2}}^x \frac{d(\psi(t) - [t])}{t} &= \int_{\frac{1}{2}}^x \frac{1}{t} d\left(\sum_{n=1}^{\infty} I(t-n)(\Lambda(n) - 1)\right) \\ &= \int_{\frac{1}{2}}^x \frac{1}{t} \sum_{n=1}^{\infty} (\Lambda(n) - 1) d\delta_n(t) \\ &= \sum_{1 \leq n \leq x} \frac{\Lambda(n) - 1}{n} \\ &= h(x). \end{aligned}$$

□

Thus, we have established that

$$\int_{\frac{1}{2}}^{\infty} \frac{d(\psi(t) - [t])}{t} = -2\gamma.$$

41. HERMITE

Hermite [42]

Hermite [43]

42. GERHARDT

Gerhardt [33, p. 196] refers to Lambert's *Architectonic*.

43. LEVI-CIVITA

Levi-Civita [57]

44. FRANEL

Franel [32] and [31]

The next theorem shows that the set of points on the unit circle that are singularities of $\sum_{n=1}^{\infty} \frac{z^n}{1-z^n}$ is dense in the unit circle. Titchmarsh [82, pp. 160–161, §4.71].

Theorem 13. For $|z| < 1$, define

$$f(z) = \sum_{n=1}^{\infty} \frac{z^n}{1-z^n}.$$

Suppose that $p > 0, q > 1$ are relatively prime integers. As $r \rightarrow 1^-$,

$$(1-r)f(re^{2\pi i/q}) \rightarrow \infty.$$

Proof. Set $z = re^{2\pi ip/q}$ and write

$$\sum_{n=1}^{\infty} \frac{z^n}{1-z^n} = \sum_{n \equiv 0 \pmod{q}} \frac{z^n}{1-z^n} + \sum_{n \not\equiv 0 \pmod{q}} \frac{z^n}{1-z^n}.$$

On the one hand,

$$\begin{aligned} (1-r) \sum_{n \equiv 0 \pmod{q}} \frac{z^n}{1-z^n} &= (1-r) \sum_{m=1}^{\infty} \frac{z^{mq}}{1-z^{mq}} \\ &= (1-r) \sum_{m=1}^{\infty} \frac{(re^{2\pi ip/q})^{mq}}{1-(re^{2\pi ip/q})^{mq}} \\ &= (1-r) \sum_{m=1}^{\infty} \frac{r^{mq}}{1-r^{mq}} \\ &= \frac{1-r}{1-r^q} \sum_{m=1}^{\infty} \frac{r^{mq}}{1+r^q+\dots+r^{(m-1)q}} \\ &= \frac{1}{1+r+\dots+r^{q-1}} \sum_{m=1}^{\infty} \frac{r^{mq}}{1+r^q+\dots+r^{(m-1)q}} \\ &\geq \frac{1}{q} \sum_{m=1}^{\infty} \frac{r^{mq}}{m} \\ &= -\frac{1}{q} \log(1-r^q) \\ &\rightarrow \infty \end{aligned}$$

as $r \rightarrow 1$.

On the other hand, for $n \not\equiv 0 \pmod{q}$ we have

$$\begin{aligned} |1-z^n|^2 &= |1-r^n e^{2\pi ipn/q}|^2 \\ &= (1-r^n e^{2\pi ipn/q})(1-r^n e^{-2\pi ipn/q}) \\ &= 1-r^n(e^{2\pi ipn/q} + e^{-2\pi ipn/q}) + r^{2n} \\ &= 1-2r^n \cos 2\pi pn/q + r^{2n} \\ &= 1-2r^n + 4r^n \sin^2 \frac{\pi pn}{q} + r^{2n} \\ &= (1-r^n)^2 + 4r^n \sin^2 \frac{\pi pn}{q}. \end{aligned}$$

So far we have not used the hypothesis that $n \equiv 0 \pmod{q}$. We use it to obtain

$$\sin \frac{\pi pn}{q} \geq \sin \frac{\pi}{q}.$$

With this we have

$$|1-z^n|^2 \geq 4r^n \sin^2 \frac{\pi}{q},$$

and therefore, as $r < 1$,

$$\begin{aligned}
(1-r) \left| \sum_{n \neq 0 \pmod{q}} \frac{z^n}{1-z^n} \right| &\leq (1-r) \sum_{n \neq 0 \pmod{q}} \frac{|z|^n}{|1-z^n|} \\
&\leq (1-r) \sum_{n \neq 0 \pmod{q}} \frac{r^n}{2r^{n/2} \sin \frac{\pi}{q}} \\
&\leq \frac{1-r}{2 \sin \frac{\pi}{q}} \sum_{n=0}^{\infty} r^{n/2} \\
&= \frac{1-r}{2 \sin \frac{\pi}{q}} \cdot \frac{1}{1-\sqrt{r}} \\
&= \frac{1+\sqrt{r}}{2 \sin \frac{\pi}{q}} \\
&< \frac{1}{\sin \frac{\pi}{q}}.
\end{aligned}$$

□

45. WIGERT

The following result is proved by Wigert [86]. Our proof follows Titchmarsh [81, p. 163, Theorem 7.15]. Cf. Landau [55].

Theorem 14. For $\lambda < \frac{1}{2}\pi$ and $N \geq 1$,

$$\sum_{n=1}^{\infty} d(n)e^{-nz} = \frac{\gamma}{z} - \frac{\log z}{z} + \frac{1}{4} - \sum_{n=0}^{N-1} \frac{B_{2n+2}^2}{(2n+2)!(2n+2)} z^{2n+1} + O(|z|^{2N})$$

as $z \rightarrow 0$ in any angle $|\arg z| \leq \lambda$.

Proof. For $\sigma > 1$, $s = \sigma + it$,

$$\zeta^2(s) = \sum_{n=1}^{\infty} \frac{d(n)}{n^s}.$$

Using this, for $\operatorname{Re} z > 0$ we have

$$\begin{aligned}
\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \Gamma(s) \zeta^2(s) z^{-s} ds &= \sum_{n=1}^{\infty} d(n) \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \Gamma(s) (nz)^{-s} ds \\
(12) \qquad \qquad \qquad &= \sum_{n=1}^{\infty} d(n) e^{-nz}.
\end{aligned}$$

Define $F(s) = \Gamma(s) \zeta^2(s) z^{-s}$. F has poles at 1, 0, and the negative odd integers. (At each negative even integer, Γ has a first order pole but ζ^2 has a second order zero.) First we determine the residue of F at 1. We use the asymptotic formula

$$\zeta(s) = \frac{1}{s-1} + \gamma + O(|s-1|), \quad s \rightarrow 1,$$

the asymptotic formula

$$\Gamma(s) = 1 - \gamma(s-1) + O(|s-1|^2), \quad s \rightarrow 1,$$

and the asymptotic formula

$$z^{-s} = \frac{1}{z} - \frac{\log z}{z}(s-1) + O(|s-1|^2), \quad s \rightarrow 1,$$

to obtain

$$\begin{aligned} \Gamma(s)\zeta^s(s)z^{-s} &= (1 - \gamma(s-1) + O(|s-1|^2)) \cdot \left(\frac{1}{(s-1)^2} + \frac{2\gamma}{s-1} + O(|s-1|^2) \right) \\ &\quad \cdot \left(\frac{1}{z} - \frac{\log z}{z}(s-1) + O(|s-1|^2) \right) \\ &= \frac{1}{z(s-1)^2} - \frac{\gamma}{z(s-1)} + \frac{2\gamma}{z(s-1)} - \frac{\log z}{z(s-1)} + O(1) \\ &= \frac{1}{z(s-1)^2} + \frac{\gamma}{z(s-1)} - \frac{\log z}{z(s-1)} + O(1). \end{aligned}$$

Hence the residue of F at 1 is

$$\frac{\gamma}{z} - \frac{\log z}{z}.$$

Now we determine the residue of F at 0. The residue of Γ at 0 is 1, and hence the residue of F at 0 is

$$1 \cdot \zeta^2(0) \cdot z^0 = \zeta^2(0) = \left(-\frac{1}{2}\right)^2 = \frac{1}{4}.$$

Finally, for $n \geq 0$ we determine the residue of F at $-(2n+1)$. The residue of Γ at $-(2n+1)$ is $\frac{(-1)^{2n+1}}{(2n+1)!}$, hence the residue of F at $-(2n+1)$ is

$$\frac{(-1)^{2n+1}}{(2n+1)!} \cdot \zeta^2(2n+1) \cdot z^{2n+1} = -\frac{B_{2n+2}^2}{(2n+2)!(2n+2)} z^{2n+1}$$

using

$$\zeta(-m) = -\frac{B_{m+1}}{m+1}, \quad m \geq 1.$$

Let $M > 0$, and let C be the rectangular path starting at $2 - iM$, then going to $2 + iM$, then going to $-2N + iM$, then going to $-2N - iM$, and then ending at $2 - iM$. By the residue theorem,

$$(13) \quad \int_C F(s)ds = 2\pi i \left(\frac{\gamma}{z} - \frac{\log z}{z} + \frac{1}{4} + \sum_{n=0}^{N-1} -\frac{B_{2n+2}^2}{(2n+2)!(2n+2)} z^{2n+1} \right).$$

Denote the right-hand side of (13) by $2\pi iR$. We have

$$\int_C F(s)ds = \int_{2-iM}^{2+iM} F(s)ds + \int_{2+iM}^{-2N+iM} F(s)ds + \int_{-2N+iM}^{-2N-iM} F(s)ds + \int_{-2N-iM}^{2-iM} F(s)ds.$$

We shall show that the second and fourth integrals tend to 0 as $M \rightarrow \infty$. For $s = \sigma + it$ with $-2N \leq \sigma \leq 2$, Stirling's formula [82, p. 151] tells us that

$$|\Gamma(s)| \sim \sqrt{2\pi} e^{-\frac{\pi}{2}|t|} |t|^{\sigma-\frac{1}{2}}, \quad |t| \rightarrow \infty.$$

As well [81, p. 95], there is some $K > 0$ such that in the half-plane $\sigma \geq -2N$,

$$\zeta(s) = O(|t|^K).$$

Also,

$$\begin{aligned} z^{-s} &= e^{-s \log z} \\ &= e^{-(\sigma+it)(\log |z|+i \arg z)} \\ &= e^{-\sigma \log |z|+t \arg z-i(\sigma \arg z+t \log |z|)}, \end{aligned}$$

and so for $|\arg z| \leq \lambda$,

$$|z^{-s}| = e^{-\sigma \log |z|+t \arg z} \leq e^{-\sigma \log |z|+\lambda |t|} = |z|^{-\sigma} e^{\lambda |t|}.$$

Therefore

$$\left| \int_{2+iM}^{-2N+iM} F(s) ds \right| \leq (2+2N) \sup_{-2N \leq \sigma \leq 2} |F(\sigma+iM)| = O(e^{-\frac{\pi}{2}M} M^{\sigma-\frac{1}{2}} M^{2K} |z|^{-\sigma} e^{\lambda M}),$$

and because $\lambda < \frac{\pi}{2}$ this tends to 0 as $M \rightarrow \infty$. Likewise,

$$\left| \int_{-2N-iM}^{2-iM} F(s) ds \right| \rightarrow 0$$

as $M \rightarrow \infty$. It follows that

$$\int_{2-i\infty}^{2+i\infty} F(s) ds + \int_{-2N+i\infty}^{-2N-i\infty} F(s) ds = 2\pi i R.$$

Hence,

$$\int_{2-i\infty}^{2+i\infty} F(s) ds = 2\pi i R + \int_{-2N-i\infty}^{-2N+i\infty} F(s) ds.$$

We bound the integral on the right-hand side. We have

$$\int_{-2N-i\infty}^{-2N+i\infty} F(s) ds = \int_{\sigma=-2N, |t| \leq 1} F(s) ds + \int_{\sigma=-2N, |t| > 1} F(s) ds.$$

The first integral satisfies

$$\left| \int_{\sigma=-2N, |t| \leq 1} F(s) ds \right| \leq \int_{\sigma=-2N, |t| \leq 1} |\Gamma(s)\zeta^2(s)| |z|^{-\sigma} e^{\lambda |t|} ds = |z|^{2N} \cdot O(1) = O(|z|^{2N}),$$

because $\Gamma(s)\zeta^2(s)$ is continuous on the path of integration. The second integral satisfies

$$\begin{aligned} \left| \int_{\sigma=-2N, |t| > 1} F(s) ds \right| &\leq \int_{\sigma=-2N, |t| > 1} e^{-\frac{\pi}{2}|t|} |t|^{\sigma-\frac{1}{2}} |t|^K |z|^{-\sigma} e^{\lambda |t|} ds \\ &= |z|^{2N} \int_{\sigma=-2N, |t| > 1} e^{-\frac{\pi}{2}|t|} |t|^{-2N-\frac{1}{2}} |t|^K e^{\lambda |t|} dt \\ &= |z|^{2N} \cdot O(1) \\ &= O(|z|^{2N}), \end{aligned}$$

because $\lambda < \frac{\pi}{2}$. This establishes

$$\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} F(s) ds = R + O(|z|^{2N}).$$

Using (12) and (13), this becomes

$$\sum_{n=1}^{\infty} d(n)e^{-nz} = \frac{\gamma}{z} - \frac{\log z}{z} + \frac{1}{4} - \sum_{n=0}^{N-1} \frac{B_{2n+2}^2}{(2n+2)!(2n+2)} z^{2n+1} + O(|z|^{-2N}),$$

completing the proof. \square

For example, as $B_2 = \frac{1}{6}, B_4 = -\frac{1}{30}, B_6 = \frac{1}{42}$, the above theorem tells us that

$$\sum_{n=1}^{\infty} d(n)e^{-nz} = \frac{\gamma}{z} - \frac{\log z}{z} + \frac{1}{4} - \frac{z}{144} - \frac{z^3}{86400} - \frac{z^5}{7620480} + O(|z|^6).$$

46. STEFFENSEN

Steffensen [75]

47. SZEGŐ

Szegő [79]

48. PÓLYA AND SZEGŐ

Pólya and Szegő [64]

49. PARTITION FUNCTION

Let

$$F(x) = \sum_{n=0}^{\infty} p(n)x^n = \prod_{n=1}^{\infty} \frac{1}{1-x^n}.$$

Taking the logarithm,

$$\log F(x) = \sum_{n=1}^{\infty} \log \frac{1}{1-x^n} = - \sum_{n=1}^{\infty} \log(1-x^n) = - \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} -\frac{(x^n)^m}{m},$$

and switching the order of summation gives

$$\log F(x) = \sum_{m=1}^{\infty} \frac{1}{m} \sum_{n=1}^{\infty} (x^m)^n = \sum_{m=1}^{\infty} \frac{1}{m} \frac{x^m}{1-x^m}.$$

On the one hand, for $0 < x < 1$ we have $mx^{m-1}(1-x) < 1-x^m$ and using this,

$$\sum_{m=1}^{\infty} \frac{1}{m} \frac{x^m}{1-x^m} < \sum_{m=1}^{\infty} \frac{1}{m} \frac{x^m}{mx^{m-1}(1-x)} = \frac{x}{1-x} \sum_{m=1}^{\infty} \frac{1}{m^2} = \frac{\pi^2}{6} \frac{x}{1-x}.$$

On the other hand, for $-1 < x < 1$ we have $1-x^m < m(1-x)$, and using this, for $0 < x < 1$ we have

$$\sum_{m=1}^{\infty} \frac{1}{m} \frac{x^m}{1-x^m} > \sum_{m=1}^{\infty} \frac{1}{m} \frac{x^m}{m(1-x)} = \frac{1}{1-x} \sum_{m=1}^{\infty} \frac{x^m}{m^2}.$$

Thus, for $0 < x < 1$,

$$\sum_{m=1}^{\infty} \frac{x^m}{m^2} < (1-x) \log F(x) < \frac{\pi^2}{6} x.$$

Taking $x \rightarrow 1^-$ gives

$$\frac{\pi^2}{6} \leq \lim_{x \rightarrow 1^-} (1-x) \log F(x) \leq \frac{\pi^2}{6},$$

i.e.,

$$\log F(x) \sim \frac{\pi^2}{6} \frac{1}{1-x}, \quad x \rightarrow 1^-.$$

See Stein and Shakarchi [76, p. 311].

50. HANSEN

Hansen [39]

51. KISELJAK

Kiseljak [45]

52. UNSORTED

In 1892, in volume VII, no. 23, p. 296 of the weekly *Naturwissenschaftliche Rundschau*, it is stated that for the year 1893, one of the six prize questions for the Belgian Academy of Sciences in Brussels is to determine the sum of the Lambert series

$$\frac{x}{1-x} + \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} + \cdots,$$

or if one cannot do this, to find a differential equation that determines the function.

Gram [35] on distribution of prime numbers.

Hardy [40]

Bohr and Cramer [1, p. 820]

Flajolet, Gourdon and Dumas [30]

REFERENCES

1. Harald Bohr and Harald Cramér, *Die neue Entwicklung der analytischen Zahlentheorie*, Encyklopädie der Mathematischen Wissenschaften mit Einschluss ihrer Anwendungen, Band II, 3. Teil, 2. Hälfte (H. Burkhardt, W. Wirtinger, R. Fricke, and E. Hilb, eds.), B. G. Teubner, Leipzig, 1923–1927, pp. 722–849.
2. Karl Bopp, *Johann Heinrich Lamberts Monatsbuch mit den zugehörigen Kommentaren, sowie mit einem Vorwort über den Stand der Lambertforschung*, Abhandlungen der Königlich Bayerischen Akademie der Wissenschaften. Mathematisch-physikalische Klasse **27** (1916), 1–84, 6. Abhandlung.
3. Peter B. Borwein, *On the irrationality of certain series*, Math. Proc. Camb. Phil. Soc. **112** (1992), 141–146.
4. V. Bouniakowsky, *Recherches sur quelques fonctions numériques*, Mémoires de l'Académie impériale des sciences de St.-Petersbourg, VIIe série **4** (1861), no. 2, 1–35.
5. Nicolas Bourbaki, *Elements of mathematics. Functions of a real variable: Elementary theory*, Springer, 2004, Translated from the French by Philip Spain.
6. T. J. I'A. Bromwich, *An introduction to the theory of infinite series*, second ed., Macmillan, London, 1959.
7. Maarten Bullynck, *Factor tables 1657–1817, with notes on the birth of number theory*, Revue d'histoire des mathématiques **16** (2010), no. 2, 133–216.
8. Heinrich Burhenne, *Ueber das Gesetz der Primzahlen*, Archiv der Mathematik und Physik **19** (1852), 442–449.
9. Eugène-Charles Catalan, *Sur la sommation de quelques séries*, Journal de Mathématiques Pures et Appliquées **7** (1842), 1–12.

10. ———, *Recherches sur quelques produits indéfinis*, Mémoires de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique **40** (1873), 1–127.
11. ———, *Mélanges mathématiques*, Mémoires de la Société Royale des Sciences de Liège, deuxième série **13** (1886), 1–404.
12. ———, *Mélanges mathématiques*, Mémoires de la Société Royale des Sciences de Liège, deuxième série **14** (1888), 1–275.
13. Augustin Cauchy, *Mémoire sur l'application du calcul des résidus au développement des produits composés d'un nombre infini de facteurs*, Comptes rendus hebdomadaires des séances de l'Académie des sciences **17** (1843), 572–581, Oeuvres complètes, série 1, tome 8, pp. 55–64.
14. ———, *Sur la réduction des rapports de factorielles réciproques aux fonctions elliptiques*, Comptes rendus hebdomadaires des séances de l'Académie des sciences **17** (1843), 825–837, Oeuvres complètes, série 1, tome 8, pp. 97–110.
15. Ernesto Cesàro, *Sur les nombres de Bernoulli et d'Euler*, Nouvelles annales de mathématiques, troisième série **5** (1886), 305–327.
16. ———, *Sur les transformations de la série de Lambert*, Nouvelles annales de mathématiques, troisième série **7** (1888), 374–382.
17. ———, *La serie di Lambert in aritmetica assintotica*, Rendiconto delle adunanze e de' lavori dell' Accademia Napolitana delle Scienze **7** (1893), 197–204.
18. ———, *Corso di analisi algebrica con introduzione al calcolo infinitesimale*, Fratelli Bocca Editori, Turin, 1894.
19. K. Chandrasekharan, *Introduction to analytic number theory*, Die Grundlehren der mathematischen Wissenschaften, vol. 148, Springer, 1968.
20. ———, *Elliptic functions*, Die Grundlehren der mathematischen Wissenschaften, vol. 281, Springer, 1985.
21. Th. Clausen, *Beitrag zur Theorie der Reihen*, J. Reine Angew. Math. **3** (1828), 92–95.
22. Henri Cohen, *Number theory, volume II: Analytic and modern tools*, Graduate Texts in Mathematics, vol. 240, Springer, 2007.
23. Maximilian Curtze, *Notes diverses sur la série de Lambert et la loi des nombres premiers*, Annali di Matematica Pura ed Applicata **1** (1867-1868), no. 1, 285–292.
24. Charles-Jean de la Vallée Poussin, *Sur la série de Lambert*, Annales de la Société Scientifique de Bruxelles **20** (1896), 56–62.
25. Gustav Lejeune Dirichlet, *Über die Bestimmung asymptotischer Gesetze in der Zahlentheorie*, Bericht über die Verhandlungen der Königlich Preussischen Akademie der Wissenschaften (1838), 13–15, Werke, Band I, pp. 351–356.
26. Sebastian Egger né Endres and Frank Steiner, *A new proof of the Voronoï summation formula*, J. Phys. A **44** (2011), no. 22, 225302.
27. G. Eisenstein, *Transformations remarquables de quelques séries*, J. Reine Angew. Math. **27** (1844), 193–197, Mathematische Werke, Band I, pp. 35–44.
28. Leonhard Euler, *Methodus generalis summandi progressionis*, Commentarii Academiae scientiarum Imperialis Petropolitanae **6** (1738), 68–97, E25, Opera omnia I.14, pp. 42–72.
29. Hans Fischer, *Dirichlet's contributions to mathematical probability theory*, Historia Math. **21** (1994), 39–63.
30. Philippe Flajolet, Xavier Gourdon, and Philippe Dumas, *Mellin transforms and asymptotics: harmonic sums*, Theoret. Comput. Sci. **144** (1995), 3–58.
31. Jérôme Franel, *Sur la théorie des séries*, Math. Ann. **52** (1899), 529–549.
32. ———, *Sur une formule utile dans la détermination de certaines valeurs asymptotiques*, Math. Ann. **51** (1899), 369–387.
33. Carl Immanuel Gerhardt, *Geschichte der Wissenschaften in Deutschland. Neuere Zeit. Siebenzehnter Band. Geschichte der Mathematik*, R. Oldenbourg, München, 1877.
34. J. W. L. Glaisher, *On the square of the series in which the coefficients are the sums of the divisors of the exponents*, Messenger of Mathematics **14** (1885), 156–163.
35. J. P. Gram, *Undersøgelser angaaende Mængden af Primtal under en given Grænse*, Det Kongelige Danske Videnskabernes Selskabs Skrifter, 6. Række, Naturvidenskabelig og Mathematisk Afdeling **2** (1881–1886), 183–308.
36. A. P. Guinand, *Functional equations and self-reciprocal functions connected with Lambert series*, Q. J. Math. **15** (1944), 11–23.
37. Siegmund Günther, *Ziele und Resultate der neueren mathematisch-historischen Forschung*, Eduard Besold, Erlangen, 1876.

38. ———, *Die Lehre von den gewöhnlichen und verallgemeinerten Hyperbelfunktionen*, Louis Nebert, Halle a. S., 1881.
39. Carl Hansen, *Note sur la sommation de la série de Lambert*, *Mathematische Annalen* **54** (1901), 604–607.
40. G. H. Hardy, *Divergent series*, second ed., AMS Chelsea Publishing, Providence, RI, 1991.
41. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, fifth ed., Oxford University Press, 1979.
42. Charles Hermite, *Sur quelques conséquences arithmétiques des formules de la théorie des fonctions elliptiques*, *Bulletin de l'Académie impériale des sciences de St.-Pétersbourg* **29** (1884), 325–352, *Œuvres*, tome IV, pp. 138–168.
43. ———, *Sur les valeurs asymptotiques de quelques fonctions numériques*, *J. Reine Angew. Math.* **99** (1886), 324–328, *Œuvres*, tome IV, pp. 209–214.
44. Carl Gustav Jacob Jacobi, *Fundamenta nova theoriae functionum ellipticarum*, Sumtibus Fratrum Borntraeger, Königsberg, 1829.
45. Marije Kiseljak, *Über Anzahlen und Summen von Teilern*, *Monatshefte für Mathematik und Physik* **28** (1917), no. 1, 133–166.
46. Georg Simon Klügel, Carl Brandan Mollweide, and Johann August Grunert (eds.), *Mathematisches Wörterbuch oder Erklärung der Begriffe, Lehrsätze, Aufgaben und Methoden der Mathematik mit den nöthigen Beweisen und literarischen Nachrichten begleitet in alphabetischer Ordnung. Erste Abtheilung. Fünfter Theil. Erster Band. I und II*, E. B. Schwickert, Leipzig, 1831.
47. Jan C. Kluyver, *On Lambert's series*, *Koninklijke Akademie van Wetenschappen te Amsterdam, Proceedings* **22** (1919), no. 4, 323–330.
48. ———, *On analytic functions defined by certain Lambert series*, *Koninklijke Akademie van Wetenschappen te Amsterdam, Proceedings* **23** (1922), no. 8, 1226–1233.
49. Konrad Knopp, *Über Lambertsche Reihen*, *J. Reine Angew. Math.* **142** (1913), 283–315.
50. W. L. Krafft, *Essai sur les nombres premiers*, *Nova Acta Academiae Scientiarum Imperialis Petropolitanae* **12** (1794), 217–245.
51. S. F. Lacroix, *Traité du calcul différentiel et du calcul intégral, tome troisième*, second ed., Veuve Courcier, Paris, 1819.
52. Edmond Laguerre, *Sur quelques théorèmes d'arithmétique*, *Bulletin de la Société Mathématique de France* **1** (1872–1873), 77–81.
53. Johann Heinrich Lambert, *Anlage zur Architectonic, oder Theorie des Einfachen und des Ersten in der philosophischen und mathematischen Erkenntniß, 2. Band*, Johann Friedrich Hartknoch, Riga, 1771.
54. Edmund Landau, *Sur la série des inverses des nombres de Fibonacci*, *Bulletin de la Société Mathématique de France* **27** (1899), 298–300.
55. ———, *Über die Wigertsche asymptotische Funktionalgleichung für die Lambertsche Reihe*, *Archiv der Mathematik und Physik*, 3. Reihe **27** (1918), 144–146, *Collected Works*, volume 7, pp. 135–137.
56. Victor-Amédée Lebesgue, *Démonstration d'une formule d'Euler, sur les diviseurs d'un nombre*, *Nouvelles annales de mathématiques* **12** (1853), 232–235.
57. Tullio Levi-Civita, *Di una espressione analitica atta a rappresentare il numero dei numeri primi compresi in un determinato intervallo*, *Atti della Reale Accademia dei Lincei, serie quinta. Rendiconti: Classe di scienze fisiche, matematiche e naturali* **4** (1895), 303–309, *Opere matematiche*, volume primo, pp. 153–158.
58. Peter Lindqvist and Jaak Peetre, *On the remainder in a series of Mertens*, *Expo. Math.* **15** (1997), no. 5, 467–478.
59. Wilhelm Lorey, *Johann Heinrich Lambert*, *Sitzungsberichte der Berliner Mathematischen Gesellschaft* **18** (1928), 2–27.
60. Friedrich Löwenhaupt (ed.), *Johann Heinrich Lambert: Leistung und Leben*, Braun & Co., Mühlhausen, 1943.
61. Franz Mertens, *Ein Beitrag zur analytischen Zahlentheorie*, *J. Reine Angew. Math.* **78** (1874), no. 1, 46–62.
62. A. F. Möbius, *Über eine besondere Art von Umkehrung der Reihen*, *J. Reine Angew. Math.* **9** (1832), no. 2, 105–123.

63. Salvatore Pincherle, *Sopra alcuni sviluppi in serie per funzioni analitiche*, Memorie della Accademia delle Scienze dell'Istituto di Bologna, serie quarta **3** (1882), 149–180, Opere scelte, vol. 1, pp. 64–91.
64. George Pólya and Gábor Szegő, *Problems and theorems in analysis, volume II*, Die Grundlehren der mathematischen Wissenschaften, vol. 216, Springer, 1976, Translated from the German by C. E. Billigheimer.
65. Franz Rogel, *Darstellung der harmonischen Reihen durch Factorenfolgen*, Archiv der Mathematik und Physik (2) **9** (1890), 297–319.
66. ———, *Darstellungen zahlentheoretischer Functionen durch trigonometrische Reihen*, Archiv der Mathematik und Physik (2) **10** (1891), 62–83.
67. Walter Rudin, *Functional analysis*, second ed., International Series in Pure and Applied Mathematics, McGraw-Hill, 1991.
68. H. F. Scherk, *Bemerkungen über die Lambertsche Reihe $\frac{x}{1-x} + \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} + \frac{x^4}{1-x^4} + \text{etc.}$* , J. Reine Angew. Math. **9** (1832), 162–168.
69. Oskar Schlömilch, *Ueber die Lambert'sche Reihe*, Zeitschrift für Mathematik und Physik **6** (1861), 407–415.
70. ———, *Extrait d'une Lettre adressée à M. Liouville par M. Schlömilch*, Journal de Mathématiques Pures et Appliquées (2) **8** (1863), 99–101.
71. ———, *Compendium der höheren Analysis, zweiter Band*, second ed., Friedrich Vieweg und Sohn, Braunschweig, 1874.
72. Francois-Joseph Servois, *Essai sur un nouveau mode d'exposition des principes du calcul différentiel*, P. Blachier-Belle, Nismes, 1814.
73. ———, *Réflexions sur les divers systèmes d'exposition des principes du calcul différentiel, et, en particulier, sur la doctrine des infiniment petits*, Annales de Mathématiques pures et appliquées **5** (1814-1815), 141–170.
74. L. A. Sohncke, *Sammlung von Aufgaben aus der Differential- und Integralrechnung*, H. W. Schmidt, Halle, 1850.
75. J. F. Steffensen, *Über Potenzreihen, im besonderen solche, deren Koeffizienten zahlentheoretische Funktionen sind*, Rendiconti del Circolo Matematico di Palermo **38** (1914), no. 1, 376–386.
76. Elias M. Stein and Rami Shakarchi, *Complex analysis*, Princeton Lectures in Analysis, vol. II, Princeton University Press, 2003.
77. M. Stern, *Beiträge zur Combinationslehre und deren Anwendung auf die Theorie der Zahlen*, J. Reine Angew. Math. **21** (1840), 177–192.
78. T.-J. Stieltjes, *Recherches sur quelques séries semi-convergentes*, Annales scientifiques de l'École Normale Supérieure, Sér. 3 **3** (1886), 201–258, Œuvres complètes, tome II, pp. 2–58.
79. Gábor Szegő, *Über Potenzreihen, deren Koeffizienten zahlentheoretische Funktionen sind*, Mathematische Zeitschrift **8** (1920), 36–51.
80. P. L. Tchebychef, *Note sur différentes séries*, Journal de Mathématiques Pures et Appliquées **16** (1851), 337–346, Œuvres, tome I, pp. 99–108.
81. E. C. Titchmarsh, *The theory of the Riemann zeta-function*, second ed., Clarendon Press, Oxford, 1986.
82. ———, *The theory of functions*, second ed., Oxford University Press, 2002.
83. Gabriele Torelli, *Sulla totalità dei numeri primi fino ad un limite assegnato*, Atti dell'Accademia delle scienze fisiche e matematiche. Sezione della Società Reale di Napoli (2) **11** (1901), no. 1, 1–222.
84. David Vernon Widder, *The Laplace transform*, Princeton Mathematical Series, vol. 6, Princeton University Press, 1946.
85. Norbert Wiener, *The Fourier integral and certain of its applications*, Cambridge University Press, 1933.
86. S. Wigert, *Sur la série de Lambert et son application à la théorie des nombres*, Acta Math. **41** (1916), 197–218.
87. A. P. Youschkevitch, *Lambert et Léonard Euler*, Colloque international et interdisciplinaire Jean-Henri Lambert, Mulhouse, 26–30 septembre 1977 (R. Oberle, A. Thill, and P. Levassort, eds.), Editions Ophrys, Paris, 1979, pp. 211–224.
88. G. Zehfuss, *Mathematische Miscellen*, Zeitschrift für Mathematik und Physik **3** (1858), 247–249.

89. Changgui Zhang, *On the modular behaviour of the infinite product $(1-x)(1-xq)(1-xq^2)(1-xq^3)\dots$* , C. R. Math. Acad. Sci. Paris **349** (2011), no. 13-14, 725–730.
90. Wadim Zudilin, *Remarks on irrationality of q -harmonic series*, Manuscripta Math. **107** (2002), no. 4, 463–477.

E-mail address: `jordan.bell@gmail.com`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA