

# Summable series and the Riemann rearrangement theorem

Jordan Bell

jordan.bell@gmail.com

Department of Mathematics, University of Toronto

May 20, 2015

## 1 Introduction

Let  $\mathbb{N}$  be the set of positive integers. A function from  $\mathbb{N}$  to a set is called a *sequence*. If  $X$  is a topological space and  $x \in X$ , a sequence  $a : \mathbb{N} \rightarrow X$  is said to *converge to  $x$*  if for every open neighborhood  $U$  of  $x$  there is some  $N_U$  such that  $n \geq N_U$  implies that  $a_n \in U$ . If there is no  $x \in X$  for which  $a$  converges to  $x$ , we say that  $a$  *diverges*.

Let  $a : \mathbb{N} \rightarrow \mathbb{R}$ . We define  $s(a) : \mathbb{N} \rightarrow \mathbb{R}$  by  $s_n(a) = \sum_{k=1}^n a_k$ . We call  $s_n(a)$  the  *$n$ th partial sum of the sequence  $a$* , and we call the sequence  $s(a)$  a *series*. If there is some  $\sigma \in \mathbb{R}$  such that  $s(a)$  converges to  $\sigma$ , we write

$$\sum_{k=1}^{\infty} a_k = \sigma.$$

## 2 Goldbach

Euler [22, §110]: “If, as is commonly the case, we take the sum of a series to be the aggregate of all of its terms, actually taken together, then there is no doubt that only infinite series that converge continually closer to some value, the more terms we actually add, can have sums”.

Euler Goldbach correspondence nos. 55, 161, 162.

## 3 Dirichlet

In 1837 Dirichlet proved that one can rearrange terms in an absolutely convergent series and not change the sum, and gave examples to show that this was not the case for conditionally convergent series.

If  $a$  is a sequence and the series  $s(|a|)$  converges, we say that the series  $s(a)$  is *absolutely convergent*. Because  $\mathbb{R}$  is a complete metric space, a series being absolutely convergent implies that it is convergent.

Dirichlet [14] and [15, p. 176, §101]  
 Elstrodt [20] and [19]

In the following theorem we prove that if a series converges absolutely, then every rearrangement of it converges to the same value. Our proof follows Landau [45, p. 157, Theorem 216].

**Theorem 1.** *If  $a$  is a sequence for which  $s(a)$  converges absolutely and*

$$\sum_{n=1}^{\infty} a_n = \sigma,$$

*then for any bijection  $\lambda : \mathbb{N} \rightarrow \mathbb{N}$ , the series  $s(a \circ \lambda)$  converges to  $\sigma$ .*

*Proof.* Let  $\epsilon > 0$ , and let  $M$  be large enough so that

$$\sum_{n=M}^{\infty} |a_n| < \epsilon.$$

Let  $r$  be large enough so that

$$\{n : 1 \leq n < M\} \subseteq \{\lambda_n : 1 \leq n \leq r\}.$$

Fix  $m \geq r$ , and let  $h : \mathbb{N} \rightarrow \mathbb{N}$  be the sequence whose terms are the elements of

$$\mathbb{N} \setminus \{\lambda_n : 1 \leq n \leq m\}$$

arranged in ascending order. If  $t + m \geq \max_{1 \leq n \leq m} \lambda_n$  then

$$\{\lambda_n : 1 \leq n \leq m\} \cup \{h_n : 1 \leq n \leq t\} = \{n : 1 \leq n \leq t + m\},$$

and hence

$$\sum_{n=1}^m a_{\lambda_n} + \sum_{n=1}^t a_{h_n} = \sum_{n=1}^{t+m} a_n.$$

Taking  $t \rightarrow \infty$ , we get

$$\sum_{n=1}^m a_{\lambda_n} + \sum_{n=1}^{\infty} a_{h_n} = \sigma;$$

the series  $s(a \circ h)$  converges because for sufficiently large  $n$ ,  $h_n = n$ . Hence, for every  $m \geq r$ ,

$$\left| \sum_{n=1}^m a_{\lambda_n} - \sigma \right| = \left| \sum_{n=1}^{\infty} a_{h_n} \right| \leq \sum_{n=1}^{\infty} |a_{h_n}| \leq \sum_{n=M}^{\infty} |a_n| < \delta,$$

which shows that  $s(a \circ \lambda)$  converges to  $\sigma$ . □

## 4 Riemann rearrangement theorem

If  $a : \mathbb{N} \rightarrow \mathbb{R}$  and  $\lambda : \mathbb{N} \rightarrow \mathbb{N}$  is a bijection, we call the sequence  $a \circ \lambda : \mathbb{N} \rightarrow \mathbb{R}$  a *rearrangement* of the sequence  $a$ .

Because  $\mathbb{N}$  is a well-ordered set, if there are at least  $n$  elements in the set  $\{k \in \mathbb{N} : a_k \geq 0\}$  then it makes sense to talk about the  $n$ th nonnegative term in the sequence  $a$ . If  $a$  were not a function from  $\mathbb{N}$  to  $\mathbb{R}$  but merely a function from a countable set to  $\mathbb{R}$ , it would not make sense to talk about the  $n$ th nonnegative term in  $a$  or the  $n$ th negative term in  $a$ .

Riemann [60, pp. 96-97]

Our proof follows Landau [45, p. 158, Theorem 217].

**Theorem 2** (Riemann rearrangement theorem). *If  $a : \mathbb{N} \rightarrow \mathbb{R}$  and  $s(a)$  converges but  $s(|a|)$  diverges, then for any nonnegative real number  $\sigma$  there is some rearrangement  $b$  of  $a$  such that  $s(b) \rightarrow \sigma$ .*

*Proof.* Define  $p, q : \mathbb{N} \rightarrow \mathbb{R}$  by

$$p_n = \frac{|a_n| + a_n}{2}, \quad q_n = \frac{|a_n| - a_n}{2}.$$

$p_n$  and  $q_n$  are nonnegative, and satisfy  $p_n - q_n = a_n$ ,  $p_n + q_n = |a_n|$ . If one of  $s(p)$  or  $s(q)$  converges and the other diverges, we obtain a contradiction from

$$s_n(a) = \sum_{k=1}^n a_k = \sum_{k=1}^n (p_k - q_k) = \sum_{k=1}^n p_k - \sum_{k=1}^n q_k = s_n(p) - s_n(q)$$

and the fact that  $s(a)$  converges. If both  $s(p)$  and  $s(q)$  converge, then we obtain a contradiction from

$$s_n(|a|) = \sum_{k=1}^n |a_k| = \sum_{k=1}^n (p_k + q_k) = \sum_{k=1}^n p_k + \sum_{k=1}^n q_k = s_n(p) + s_n(q)$$

and the fact that  $s(|a|)$  diverges. Therefore, both  $s(p)$  and  $s(q)$  diverge.

Because  $s(a)$  converges and  $s(|a|)$  diverges, there are infinitely many  $n$  with  $a_n > 0$  and there are infinitely many  $n$  with  $a_n < 0$ . Let  $P_n$  be the  $n$ th nonnegative term in the sequence  $a$ , and let  $Q_n$  be the absolute value of the  $n$ th negative term in the sequence  $a$ . The fact that  $s(p)$  diverges implies that  $s(P)$  diverges, and the fact that  $s(q)$  diverges implies that  $s(Q)$  diverges.

Let  $\sigma \geq 0$ . We define sequences  $\mu, \nu : \mathbb{N} \rightarrow \mathbb{N}$  by induction as follows. Let  $\mu_1$  be the least element of  $\mathbb{N}$  such that

$$s_{\mu_1}(P) > \sigma,$$

and with  $\mu_1$  chosen, let  $\nu_1$  be the least element of  $\mathbb{N}$  such that

$$s_{\mu_1}(P) - s_{\nu_1}(Q) < \sigma.$$

Let  $m_2$  be the least element of  $\mathbb{N}$  such that

$$s_{\mu_2}(P) - s_{\nu_1}(Q) > \sigma,$$

and with  $\mu_2$  chosen, let  $\nu_2$  be the least element of  $\mathbb{N}$  such that

$$s_{\mu_2}(P) - s_{\nu_2}(Q) < \sigma.$$

It is straightforward to check that  $\mu_2 > \mu_1$  and  $\nu_2 > \nu_1$ .

Suppose that  $\mu_1, \dots, \mu_n$  and  $\nu_1, \dots, \nu_n$  have been chosen, that  $\mu_n$  is the least element of  $\mathbb{N}$  such that

$$s_{\mu_n}(P) - s_{\nu_{n-1}}(Q) > \sigma,$$

that  $\nu_n$  is the least element of  $\mathbb{N}$  such that

$$s_{\mu_n}(P) - s_{\nu_n}(Q) < \sigma$$

and that  $\mu_n > \mu_{n-1}$  and  $\nu_n > \nu_{n-1}$ . Let  $\mu_{n+1}$  be the least element of  $\mathbb{N}$  such that

$$s_{\mu_{n+1}}(P) - s_{\nu_n}(Q) > \sigma,$$

and with  $\mu_{n+1}$  chosen, let  $\nu_{n+1}$  be the least element of  $\mathbb{N}$  such that

$$s_{\mu_{n+1}}(P) - s_{\nu_{n+1}}(Q) < \sigma.$$

It is straightforward to check that  $\mu_{n+1} > \mu_n$  and  $\nu_{n+1} > \nu_n$ .

Define  $b : \mathbb{N} \rightarrow \mathbb{R}$  by taking  $b_n$  to be the  $n$ th term in

$$P_1, \dots, P_{\mu_1}, -Q_1, \dots, -Q_{\nu_1}, P_{\mu_1+1}, \dots, P_{\mu_2}, -Q_{\nu_1+1}, \dots, -Q_{\nu_2}, \dots,$$

which, because the sequences  $\mu$  and  $\nu$  are strictly increasing, is a rearrangement of the sequence  $a$ . □

## 5 Symmetry

Don't use order where it is accidental.

## 6 Nets

A *directed set* is a set  $D$  and a binary relation  $\preceq$  satisfying

- if  $m, n, p \in D$ ,  $m \preceq n$ , and  $n \preceq p$ , then  $m \preceq p$
- if  $m \in D$ , then  $m \preceq m$
- if  $m, n \in D$ , then there is some  $p \in D$  such that  $m \preceq p$  and  $n \preceq p$ .

For example, let  $A$  be a set, let  $D$  be the set of all subsets of  $A$ , and say that  $F \preceq G$  when  $F \subseteq G$ . Check that  $(D, \preceq)$  is a directed set: for  $F, G \in D$ , we have  $F \cup G \in D$ , and  $F \cup G$  is an upper bound for both  $F$  and  $G$ .

A *net* is a function from a directed set  $(D, \preceq)$  to a set  $X$ . Let  $(X, \tau)$  be a topological space, let  $S : (D, \preceq) \rightarrow (X, \tau)$  be a net, and let  $x \in X$ . We say that  $S$  *converges to  $x$*  if for every  $U \in \tau$  with  $x \in U$  there is some  $N_U \in D$  such that  $N_U \preceq i$  implies that  $S(i) \in U$ . One proves that a topological space is Hausdorff if and only if every net in this space converges to at most one point [41, p. 67, Theorem 3].

A net  $S : (D, \preceq) \rightarrow \mathbb{R}$  is said to be *increasing* if  $m \preceq n$  implies that  $S(m) \leq S(n)$ .

**Lemma 3.** *If  $S : (D, \preceq) \rightarrow \mathbb{R}$  is an increasing net and the range  $R$  of  $S$  has an upper bound, then  $S$  converges to the supremum of  $R$ .*

*Proof.* Because  $R$  is a subset of  $\mathbb{R}$  that has an upper bound, it has a supremum, call it  $\sigma$ . To say that  $\sigma$  is the supremum of  $R$  means that for all  $r \in R$  we have  $r \leq \sigma$  ( $\sigma$  is an upper bound) and that for all  $\epsilon > 0$  there is some  $r_\epsilon \in R$  with  $\sigma - \epsilon < r_\epsilon$  (nothing less than  $\sigma$  is an upper bound). Take  $\epsilon > 0$ . There is some  $r_\epsilon \in R$  with  $\sigma - \epsilon < r_\epsilon$ . As  $r_\epsilon \in R$ , there is some  $n_\epsilon \in D$  with  $S(n_\epsilon) = r_\epsilon$ . If  $n_\epsilon \preceq n$ , then because  $S$  is increasing,  $S(n_\epsilon) \leq S(n)$ , and hence

$$\sigma - \epsilon < r_\epsilon = S(n_\epsilon) \leq S(n).$$

But  $S(n) \in R$ , so  $S(n) \leq \sigma$ . Hence  $n_\epsilon \preceq n$  implies that  $|S(n) - \sigma| < \epsilon$ , showing that  $S$  converges to  $\sigma$ .  $\square$

## 7 Unordered sums

Let  $A$  be a set, and let  $\mathcal{P}_0(A)$  be the set of all finite subsets of  $A$ . Check that  $(\mathcal{P}_0(A), \subseteq)$  is a directed set: if  $F, G \in \mathcal{P}_0(A)$  then  $F \cup G \in \mathcal{P}_0(A)$  and  $F \cup G$  is an upper bound for both  $F$  and  $G$ . Let  $f : A \rightarrow \mathbb{R}$  be a function, and define  $S_f : \mathcal{P}_0(A) \rightarrow \mathbb{R}$  by

$$S_f(F) = \sum_{a \in F} f(a), \quad F \in \mathcal{P}_0(A).$$

If the net  $S_f$  converges, we say that the function  $f$  is *summable*, and we call the element of  $\mathbb{R}$  to which  $S_f$  converges the *unordered sum of  $f$* , denoted by

$$\sum_{a \in A} f(a).$$

If  $B$  is a subset of  $A$ , we say that  $f$  is *summable over  $B$*  if the restriction of  $f$  to  $B$  is summable. If  $f_B$  is the restriction of  $f$  to  $B$  and  $f$  is summable over  $B$  (i.e.  $f_B$  is summable), by

$$\sum_{a \in B} f(a)$$

we mean

$$\sum_{a \in B} f_B(a).$$

**Lemma 4.** *Suppose that  $f, g : A \rightarrow \mathbb{R}$  are functions and  $\alpha, \beta \in \mathbb{R}$ . If  $f$  and  $g$  are summable, then  $\alpha f + \beta g$  is summable and*

$$\sum_{a \in A} (\alpha f(a) + \beta g(a)) = \alpha \sum_{a \in A} f(a) + \beta \sum_{a \in A} g(a).$$

*Proof.* Let  $\sigma_1 = \sum_{a \in A} f(a)$  and  $\sigma_2 = \sum_{a \in A} g(a)$ , and set  $h = \alpha f + \beta g$ . For  $\epsilon > 0$ , there is some  $F_\epsilon \in \mathcal{P}_0(A)$  such that  $F_\epsilon \subseteq F \in \mathcal{P}_0(A)$  implies that  $|S_f(F) - \sigma_1| < \epsilon$ , and there is some  $G_\epsilon \in \mathcal{P}_0(A)$  such that  $G_\epsilon \subseteq G \in \mathcal{P}_0(A)$  implies that  $|S_g(G) - \sigma_2| < \epsilon$ . Let  $H_\epsilon = F_\epsilon \cup G_\epsilon \in \mathcal{P}_0(A)$ . If  $H_\epsilon \subseteq H \in \mathcal{P}_0(A)$ , then, as  $F_\epsilon \subseteq H$  and  $G_\epsilon \subseteq H$ ,

$$\begin{aligned} |S_h(H) - (\alpha\sigma_1 + \beta\sigma_2)| &= \left| \sum_{a \in H} (\alpha f(a) + \beta g(a)) - \alpha\sigma_1 - \beta\sigma_2 \right| \\ &= |\alpha S_f(H) + \beta S_g(H) - \alpha\sigma_1 - \beta\sigma_2| \\ &\leq |\alpha| |S_f(H) - \sigma_1| + |\beta| |S_g(H) - \sigma_2| \\ &\leq |\alpha|\epsilon + |\beta|\epsilon; \end{aligned}$$

we write  $\leq$  rather than  $<$  in the last inequality to cover the case where  $\alpha = \beta = 0$ . It follows that  $S_h$  converges to  $\alpha\sigma_1 + \beta\sigma_2$ .  $\square$

The following lemma is simple to prove and ought to be true, but should not be called obvious. For example, the Cesàro sum of the sequence  $1, -1, 1, -1, \dots$  is  $\frac{1}{2}$ , while the Cesàro sum of the sequence  $1, -1, 0, 1, -1, 0, \dots$  is  $\frac{1}{3}$ .

**Lemma 5.** *If  $f : A \rightarrow \mathbb{R}$  is summable, then for any set  $C$  that contains  $A$ , the function  $g : C \rightarrow \mathbb{R}$  defined by*

$$g(c) = \begin{cases} f(c) & c \in A \\ 0 & \text{otherwise} \end{cases}$$

*is summable, and*

$$\sum_{a \in A} f(a) = \sum_{c \in C} g(c).$$

*Proof.* Let  $\sigma = \sum_{a \in A} f(a)$ . For  $\epsilon > 0$ , there is some  $F_\epsilon \in \mathcal{P}_0(A)$  such that  $F_\epsilon \subseteq F \in \mathcal{P}_0(A)$  implies that  $|S_f(F) - \sigma| < \epsilon$ . If  $F_\epsilon \subseteq H \in \mathcal{P}_0(C)$ , then, as

$F_\epsilon \subseteq H \cap A \in \mathcal{P}_0(A)$ ,

$$\begin{aligned}
|S_g(H) - \sigma| &= \left| \sum_{c \in H} g(c) - \sigma \right| \\
&= \left| \sum_{c \in H \cap A} g(c) + \sum_{c \in H \setminus A} g(c) - \sigma \right| \\
&= \left| \sum_{a \in H \cap A} f(a) + \sum_{c \in H \setminus A} 0 - \sigma \right| \\
&= |S_f(H \cap A) - \sigma| \\
&< \epsilon.
\end{aligned}$$

This shows that  $S_g$  converges to  $\sigma$ . □

The previous two lemmas are useful, and also convince us that unordered summation works similarly to finite sums. We now establish conditions under which a function is summable.

**Lemma 6.** *If  $f : A \rightarrow \mathbb{R}$  is nonnegative and there is some  $M \in \mathbb{R}$  such that  $F \in \mathcal{P}_0(A)$  for all  $S_f(F) \leq M$ , then  $f$  is summable. If  $f : A \rightarrow \mathbb{R}$  is nonnegative and summable, then  $S_f(F) \leq \sum_{a \in A} f(a)$  for all  $F \in \mathcal{P}_0(A)$ .*

*Proof.* Suppose there is some  $M \in \mathbb{R}$  such that if  $F \in \mathcal{P}_0(A)$  then  $S_f(F) \leq M$ . That is,  $M$  is an upper bound for the range of  $S_f$ . Because  $f$  is nonnegative, the net  $S_f$  is increasing. We apply Lemma 3, which tells us that  $S_f$  converges to the supremum of its range. That  $S_f$  converges means that  $f$  is summable.

Suppose that  $f$  is summable, and let  $\sigma = \sum_{a \in A} f(a)$ . Suppose by contradiction that there is some  $F_0 \in \mathcal{P}_0(A)$  such that  $S_f(F_0) > \sigma$ , and let  $\epsilon = S_f(F_0) - \sigma$ . Then there is some  $F_\epsilon \in \mathcal{P}_0(A)$  such that  $F_\epsilon \subseteq F_0 \in \mathcal{P}_0(A)$  implies that  $|S_f(F_\epsilon) - \sigma| < \epsilon$ . As  $F_\epsilon \subseteq F_0 \cup F_\epsilon \in \mathcal{P}_0(A)$ , we have  $|S_f(F_0 \cup F_\epsilon) - \sigma| < \epsilon$ , and hence

$$S_f(F_0 \cup F_\epsilon) < \sigma + \epsilon = S_f(F_0).$$

But  $F_0$  is contained in  $F_0 \cup F_\epsilon$  and  $f$  is nonnegative, so

$$S_f(F_0) \leq S_f(F_0 \cup F_\epsilon),$$

which gives  $S_f(F_0) < S_f(F_0)$ , a contradiction. Therefore, there is no  $F \in \mathcal{P}_0(A)$  for which  $S_f(F) > \sigma$ . □

**Lemma 7.** *Suppose that  $f : A \rightarrow \mathbb{R}$  is a function and that  $A_+ = \{a \in A : f(a) \geq 0\}$  and  $A_- = \{a \in A : f(a) \leq 0\}$ . Then,  $f$  is summable if and only if  $f$  is summable over both  $A_+$  and  $A_-$ . If  $f$  is summable, then*

$$\sum_{a \in A} f(a) = \sum_{a \in A_+} f(a) + \sum_{a \in A_-} f(a).$$

*Proof.* Suppose that  $f$  is summable. Because  $f$  is summable, there is some  $E \in \mathcal{P}_0(A)$  such that  $E \subseteq F \in \mathcal{P}_0(A)$  implies that  $|S_f(F) - \sigma| < 1$ . Define

$$E_+ = \{a \in E : f(a) \geq 0\} \in \mathcal{P}_0(A_+), \quad E_- = \{a \in E : f(a) \leq 0\} \in \mathcal{P}_0(A_-).$$

If  $G \in \mathcal{P}_0(A_+)$  then  $E \subseteq G \cup E \in \mathcal{P}_0(A)$ , and hence  $|S_f(G \cup E) - \sigma| < 1$ . We have

$$S_{f_+}(G) = \sum_{a \in G} f(a) \leq \sum_{a \in G \cup E_+} f(a) = \sum_{a \in G \cup E} f(a) - \sum_{a \in E_-} f(a),$$

and hence

$$S_{f_+}(G) \leq S_f(G \cup E) - S_f(E_-) < \sigma + 1 - S_f(E_-).$$

That is,  $\sigma + 1 - S_f(E_-)$  is an upper bound for the range of  $S_{f_+}$ . The net  $S_{f_+}$  is increasing, hence applying Lemma 3 we get that  $S_{f_+}$  converges. That is,  $f_+$  is summable. If  $H \in \mathcal{P}_0(A_-)$ , then  $E \subseteq H \cup E \in \mathcal{P}_0(A)$ , and hence  $|S_f(H \cup E) - \sigma| < 1$ . We have

$$S_{f_-}(H) = \sum_{a \in H} f(a) \geq \sum_{a \in H \cup E_-} f(a) = \sum_{a \in H \cup E} f(a) - \sum_{a \in E_+} f(a),$$

and then

$$S_{f_-}(H) \geq S_f(H \cup E) - S_f(E_+) > \sigma - 1 - S_f(E_+),$$

showing that  $-\sigma + 1 + S_f(E_+)$  is an upper bound for the net  $-S_{f_-}$ . As  $-S_{f_-}$  is increasing, by Lemma 3 it converges, and it follows that  $S_{f_-}$  converges. That is,  $f_-$  is summable.

Suppose that  $f$  is summable over both  $A_+$  and  $A_-$ . Let  $f_+$  be the restriction of  $f$  to  $A_+$  and let  $f_-$  be the restriction of  $f$  to  $A_-$ , and define  $g_+, g_- : A \rightarrow \mathbb{R}$  by

$$g_+(a) = \begin{cases} f(a) & a \in A_+ \\ 0 & a \in A_- \end{cases}, \quad g_-(a) = \begin{cases} 0 & a \in A_+ \\ f(a) & a \in A_- \end{cases}.$$

By Lemma 5,  $f_+$  being summable implies that  $g_+$  is summable, with

$$\sum_{a \in A_+} f_+(a) = \sum_{a \in A} g_+(a),$$

and  $f_-$  being summable implies that  $g_-$  is summable, with

$$\sum_{a \in A_-} f_-(a) = \sum_{a \in A} g_-(a).$$

But  $f = g_+ + g_-$ , so by Lemma 4 we get that  $f$  is summable, with

$$\sum_{a \in A} f(a) = \sum_{a \in A} g_+(a) + \sum_{a \in A} g_-(a) = \sum_{a \in A_+} f_+(a) + \sum_{a \in A_-} f_-(a).$$

□



If  $f : A \rightarrow \mathbb{R}$  is a function, we define  $|f| : A \rightarrow \mathbb{R}$  by  $|f|(a) = |f(a)|$ .

**Theorem 8.** *If  $f : A \rightarrow \mathbb{R}$  is a function, then  $f$  is summable if and only if  $|f|$  is summable.*

*Proof.* Let  $A_+ = \{a \in A : f(a) \geq 0\}$  and  $A_- = \{a \in A : f(a) \leq 0\}$ , and let  $f_+$  and  $f_-$  be the restrictions of  $f$  to  $A_+$  and  $A_-$  respectively. Suppose that  $f$  is summable. Then by Lemma 7 we get that  $f_+$  is summable and  $f_-$  is summable. Let  $F \in \mathcal{P}_0(A)$  and write  $F_+ = \{a \in F : f(a) \geq 0\}$ ,  $F_- = \{a \in F : f(a) \leq 0\}$ . We have

$$S_{|f|}(F) = \sum_{a \in F} |f(a)| = \sum_{a \in F_+} f(a) - \sum_{a \in F_-} f(a) = S_{f_+}(F_+) - S_{f_-}(F_-).$$

But by Lemma 6, because the net  $S_{f_+}$  is increasing we have  $S_{f_+}(F_+) \leq \sum_{a \in A_+} f_+(a)$ , and because the net  $-S_{f_-}$  is increasing we have  $-S_{f_-}(F_-) \leq -\sum_{a \in A_-} f_-(a)$ . Therefore,  $\sum_{a \in A_+} f_+(a) - \sum_{a \in A_-} f_-(a)$  is an upper bound for the range of  $S_{|f|}$ . Moreover,  $S_{|f|}$  is increasing, so by Lemma 6 it follows that  $S_{|f|}$  converges, i.e. that  $|f|$  is summable.

Suppose that  $|f|$  is summable. By Lemma 6, for any  $F \in \mathcal{P}_0(A_+)$  we have

$$S_{f_+}(F) = S_{|f|}(F) \leq \sum_{a \in A} |f(a)|,$$

i.e.,  $\sum_{a \in A} |f(a)|$  is an upper bound for the range of  $S_{f_+}$ . As  $S_{f_+}$  is increasing, by Lemma 6 it follows that  $S_{f_+}$  converges, i.e., that  $f_+$  is summable. Because  $-S_{f_-}$  is increasing, we likewise get that  $-S_{f_-}$  converges and hence that  $S_{f_-}$  converges, i.e. that  $f_-$  is summable. Now applying Lemma 7, we get that  $f$  is summable.  $\square$

**Theorem 9.** *If  $f : A \rightarrow \mathbb{R}$  is summable, then  $\{a \in A : f(a) \neq 0\}$  is countable.*

*Proof.* Suppose by contradiction that  $\{a \in A : f(a) \neq 0\}$  is uncountable. We have

$$\{a \in A : f(a) \neq 0\} = \{a \in A : |f(a)| > 0\} = \bigcup_{n \in \mathbb{N}} \left\{ a \in A : |f(a)| \geq \frac{1}{n} \right\}.$$

Since this is a countable union, there is some  $n \in \mathbb{N}$  such that  $\{a \in A : |f(a)| \geq \frac{1}{n}\}$  is uncountable; in particular, this set is infinite. Because  $f$  is summable, by Theorem 8 we have that  $|f|$  is summable, with unordered sum  $\sigma$ . Hence, there is some  $F_1 \in \mathcal{P}_0(A)$  such that  $F_1 \subseteq F \in \mathcal{P}_0(A)$  implies that  $|S_{|f|}(F) - \sigma| < 1$ . Let  $F$  be a finite subset of  $\{a \in A : |f(a)| \geq \frac{1}{n}\}$  with at least  $n(\sigma + 1)$  elements. Then

$$S_{|f|}(F \cup F_1) = \sum_{a \in F \cup F_1} |f(a)| \geq \sum_{a \in F} |f(a)| \geq n(\sigma + 1) \cdot \frac{1}{n} = \sigma + 1.$$

But  $F_1 \subseteq F \cup F_1 \in \mathcal{P}_0(A)$ , so  $S_{|f|}(F \cup F_1) < \sigma + 1$ , a contradiction. Therefore,  $\{a \in A : f(a) \neq 0\}$  is countable.  $\square$

## 8 References

McArthur [52]  
Schaefer [64, p. 120]  
Roytvarf [63, p. 282]  
McShane [53]  
Diestel, Jarchow and Tonge [13]  
Remmert [59, p. 29]  
Sorenson [67]  
Lattice sums [18]  
Kadets and Kadets [40]  
Manning [50]  
Bottazini [2]  
Boyer [4]  
Weil [74]  
Smithies [66]  
Dugac [16]  
Grattan-Guinness [31] and [33] and [32] and [30]  
Whiteside [75]  
Schaefer [65]  
Cauchy [5]  
Polya [57]  
Lakatos [44]  
Krantz [43]  
Cunha [12]  
Youshkevitch [77]  
Tweddle [70] and [71]  
Bromwich [6, p. 74, Art. 28]  
Laugwitz [46], [47], [48]  
Tucciarone [69]  
Fraser [27]  
Cowen [11]  
Spence [68]  
Jahnke [38]  
Epple [21]  
Mascré [51]  
Rosenthal [61]  
Freniche [28]  
Goursat [29, p. 348]

## 9 Probability

Baker [1]  
Nathan [55]  
Nover and Harris [56]

Colyvan [10]  
 Liouville [49, pp. 74–75]  
 Chrystal [9, p. 118]  
 Jordan [39, p. 277, Theorem 291]  
 Cayley [8, a]  
 Harkness and Morley [35, p. 66]  
 Hofmann [37]  
 Ferreirós [26]  
 Ferraro [25] and [23] and [24]  
 Brouncker [7]  
 Roy [62]  
 Wallis [72]  
 Bourbaki [3, p. 261, chapter III, §5.1]  
 Pringsheim [58]  
 Dutka [17]  
 Grünbaum [34]  
 Hinton and Martin [36]  
 Gersonides [42]  
 Watling [73]  
 Moore [54]  
 Wojtaszczyk [76, Chapter 7]

## References

- [1] Alan Baker, *Putting expectations in order*, *Philosophy of Science* **74** (2007), no. 5, 692–700.
- [2] Umberto Bottazzini, *The higher calculus: A history of real and complex analysis from Euler to Weierstrass*, Springer, 1986, Translated from the Italian by Warren Van Egmond.
- [3] Nicolas Bourbaki, *General topology, chapters 1–4*, *Elements of Mathematics*, Springer, 1989.
- [4] Carl B. Boyer, *The history of the calculus and its conceptual development*, Dover Publications, New York, 1959.
- [5] Robert E. Bradley and C. Edward Sandifer, *Cauchy’s Cours d’analyse: an annotated translation*, *Sources and Studies in the History of Mathematical and Physical Sciences*, Springer, 2009.
- [6] T. J. F. A. Bromwich, *An introduction to the theory of infinite series*, second ed., Macmillan and Co., London, 1931.
- [7] William Brouncker, *The squaring of the hyperbola, by an infinite series of rational numbers, together with its demonstration, by that eminent mathematician, the right honourable the Lord Viscount Brouncker*, *Phil. Trans.* **3** (1668), 645–649.

- [8] Arthur Cayley, *Series*, Encyclopædia Britannica, vol. XXI, Charles Scribner's Sons, New York, ninth ed., 1886, Collected Mathematical Papers, vol. XI, pp. 617–627, pp. 677–682.
- [9] George Chrystal, *Algebra: An elementary text-book for the higher classes of secondary schools and for colleges, part II*, Adam and Charles Black, Edinburgh, 1889.
- [10] Mark Colyvan, *No expectations*, *Mind* **115** (2006), no. 459, 695–702.
- [11] C. C. Cowen, K. R. Davidson, and R. P. Kaufman, *Rearranging the alternating harmonic series*, *Amer. Math. Monthly* **87** (1980), no. 10, 817–819.
- [12] A. J. Franco de Oliveira, *Anastácio da Cunha and the concept of convergent series*, *Arch. Hist. Exact Sci.* **39** (1988), no. 1, 1–12.
- [13] Joe Diestel, Hans Jarchow, and Andrew Tonge, *Absolutely summing operators*, Cambridge Studies in Advanced Mathematics, vol. 43, Cambridge University Press, 1995.
- [14] Gustav Lejeune Dirichlet, *Beweis des Satzes, dass jede unbegrenzte arithmetische Progression, deren erstes Glied und Differenz ganze Zahlen ohne gemeinschaftlichen Factor sind, unendlich viele Primzahlen enthält*, Abhandlungen der Königlich Akademie der Wissenschaften zu Berlin (1837), 45–81, Werke, vol. I, pp. 313–342.
- [15] P. G. L. Dirichlet, *Lectures on number theory*, History of Mathematics, vol. 16, American Mathematical Society, Providence, RI, 1999, Supplements by R. Dedekind, translated from the German by John Stilwell.
- [16] Pierre Dugac, *Éléments d'analyse de Karl Weierstrass*, *Arch. History Exact Sci.* **10** (1973), no. 1/2, 41–176.
- [17] Jacques Dutka, *On the St. Petersburg paradox*, *Arch. Hist. Exact Sci.* **39** (1988), no. 1, 13–39.
- [18] Ulrich Eckhardt, *Gittersummen der Festkörperphysik. Wie sinnvoll ist die Sprache des Unendlichen für die Dinge des alltäglichen Lebens?*, *Mitt. Math. Ges. Hamburg* **29** (2010), 71–96.
- [19] Jürgen Elstrodt, *Annotationes historicae*, *Mitt. Dtsch. Math.-Ver.* (1998), no. 1, 23–24.
- [20] Jürgen Elstrodt, *The life and work of Gustav Lejeune Dirichlet (1805–1859)*, *Analytic Number Theory: A Tribute to Gauss and Dirichlet* (William Duke and Yuri Tschinkel, eds.), Clay Mathematics Proceedings, vol. 7, American Mathematical Society, Providence, RI, 2005, pp. 1–38.

- [21] Moritz Epple, *The end of the science of quantity: foundations of analysis, 1860–1910*, A History of Analysis (Hans Niels Jahnke, ed.), History of Mathematics, vol. 24, American Mathematical Society, Providence, RI, 2003, pp. 291–324.
- [22] Leonhard Euler, *Institutiones calculi differentialis, vol. I*, Academia Scientiarum Imperialis Petropolitanae, St. Petersburg, 1755, E212, Opera omnia I.10.
- [23] Giovanni Ferraro, *The first modern definition of the sum of a divergent series: an aspect of the rise of 20th century mathematics*, Arch. Hist. Exact Sci. **54** (1999), no. 2, 101–135.
- [24] ———, *True and fictitious quantities in Leibniz’s theory of series*, Studia Leibnitiana **32** (2000), no. 1, 43–67.
- [25] ———, *The rise and development of the theory of series up to the early 1820s*, Sources and Studies in the History of Mathematics and Physical Sciences, Springer, 2008.
- [26] José Ferreirós, *Labyrinth of thought: A history of set theory and its role in modern mathematics*, second ed., Birkhäuser, 2007.
- [27] Craig G. Fraser, *The calculus as algebraic analysis: some observations on mathematical analysis in the 18th century*, Arch. Hist. Exact Sci. **39** (1989), no. 4, 317–335.
- [28] Francisco J. Freniche, *On Riemann’s rearrangement theorem for the alternating harmonic series*, Amer. Math. Monthly **117** (2010), no. 5, 442–448.
- [29] Édouard Goursat, *A course in mathematical analysis, vol. I*, Ginn and Company, Boston, 1904, Translated from the French by Earle Raymond Hedrick.
- [30] Ivor Grattan-Guinness, *Bolzano, Cauchy and the “new analysis” of the early nineteenth century*, Arch. Hist. Exact Sci. **6** (1970), no. 5, 372–400.
- [31] ———, *The development of the foundations of mathematical analysis from Euler to Riemann*, MIT Press, 1970.
- [32] ———, *The emergence of mathematical analysis and its foundational progress, 1780–1880*, From the Calculus to Set Theory, 1630–1910 (Ivor Grattan-Guinness, ed.), Princeton University Press, 1980, pp. 94–148.
- [33] ———, *Thus it mysteriously appears: impressions of Laplace’s use of series*, Rechnen mit dem Unendlichen: Beiträge zur Entwicklung eines kontroversen Gegenstandes (Detlef D. Spalt, ed.), Birkhäuser, 1990, pp. 95–102.
- [34] Adolf Grünbaum, *Can an infinitude of operations be performed in a finite time?*, Br. J. Philos. Sci. **20** (1969), no. 3, 203–218.

- [35] James Harkness and Frank Morley, *A treatise on the theory of functions*, Macmillan, London, 1893.
- [36] J. M. Hinton and C. B. Martin, *Achilles and the tortoise*, *Analysis* **14** (1954), no. 3, 56–68.
- [37] Joseph Ehrenfried Hofmann, *On the discovery of the logarithmic series and its development in England up to Cotes*, *National Mathematics Magazine* **14** (1939), no. 1, 37–45.
- [38] Hans Niels Jahnke, *Algebraic analysis in the 18th century*, *A History of Analysis* (Hans Niels Jahnke, ed.), *History of Mathematics*, vol. 24, American Mathematical Society, Providence, RI, 2003, pp. 105–136.
- [39] M. C. Jordan, *Cours d’analyse de l’école Polytechnique, tome premier*, second ed., Gauthier-Villars et fils, Paris, 1893.
- [40] Mikhail I. Kadets and Vladimir M. Kadets, *Series in Banach spaces: Conditional and unconditional convergence*, *Operator Theory: Advances and Applications*, vol. 94, Birkhäuser, 1997, Translated from the Russian by Andrei Iacob.
- [41] John L. Kelley, *General topology*, *University Series in Higher Mathematics*, D. Van Nostrand Company, 1955.
- [42] George Kohler, *Medieval infinities in mathematics and the contribution of Gersonides*, *History of Philosophy Quarterly* **23** (2006), no. 2, 95–116.
- [43] Steven G. Krantz and Jeffery D. McNeal, *Creating more convergent series*, *Amer. Math. Monthly* **111** (2004), no. 1, 32–38.
- [44] Imre Lakatos, *Proofs and refutations*, Cambridge University Press, 1976.
- [45] Edmund Landau, *Differential and integral calculus*, Chelsea Publishing Company, New York, 1951, Translated from the German by Melvin Hausner and Martin Davis.
- [46] Detlef Laugwitz, *Definite values of infinite sums: Aspects of the foundations of infinitesimal analysis around 1820*, *Arch. Hist. Exact Sci.* **39** (1989), no. 3, 195–245.
- [47] ———, *Controversies about numbers and functions*, *The Growth of Mathematical Knowledge* (Emily Grosholz and Herbert Breger, eds.), *Sythese Library*, vol. 289, Kluwer, 2000, pp. 177–198.
- [48] ———, *Bernhard Riemann 1826–1866: Turning points in the conception of mathematics*, Birkhäuser, 2008, Translated from the German by Abe Shenitzer.

- [49] Jesper Lützen, *Joseph Liouville 1809–1882: master of pure and applied mathematics*, Studies in the History of Mathematics and Physical Sciences, vol. 15, Springer, 1990.
- [50] Kenneth R. Manning, *The emergence of the Weierstrassian approach to complex analysis*, Arch. Hist. Exact Sci. **14** (1975), no. 4, 297–383.
- [51] David Mascré, *1867 Bernhard Riemann, posthumous thesis on the representation of functions by trigonometric series*, Landmark Writings in Western Mathematics 1640–1940 (Ivor Grattan-Guinness, ed.), Elsevier, 2005, pp. 491–505.
- [52] C. W. McArthur, *Series with sums invariant under rearrangement*, Amer. Math. Monthly **75** (1968), 729–731.
- [53] E. J. McShane, *Partial orderings and Moore-Smith limits*, Amer. Math. Monthly **59** (1952), no. 1, 1–11.
- [54] C. N. Moore, *Summability of series*, Amer. Math. Monthly **39** (1932), no. 2, 62–71.
- [55] Amos Nathan, *False expectations*, Philosophy of Science **51** (1984), no. 1, 128–136.
- [56] Harris Nover and Alan Hájek, *Vexing expectations*, Mind **113** (2004), no. 450, 237–249.
- [57] George Pólya and Gábor Szegő, *Problems and theorems in analysis, volume I*, Die Grundlehren der mathematischen Wissenschaften, vol. 193, Springer, 1972, Translated from the German by D. Aeppli.
- [58] Alfred Pringsheim, *Irrationalzahlen und Konvergenz unendlicher Prozesse*, Enzyklopädie der Mathematischen Wissenschaften mit Einschluss ihrer Anwendungen. Band I, 1. Teil (Wilhelm Franz Meyer, ed.), B. G. Teubner, Leipzig, 1898–1904, pp. 47–146.
- [59] Reinhold Remmert, *Theory of complex functions*, Graduate Texts in Mathematics, vol. 122, Springer, 1991, Translated from the German by Robert B. Burckel.
- [60] Bernhard Riemann, *Ueber die Darstellbarkeit einer Function durch eine trigonometrische Reihe*, Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen **13** (1866/1867), 87–132, Gesammelte mathematische Werke, second ed., vol. II, 227–271.
- [61] Peter Rosenthal, *The remarkable theorem of Lévy and Steinitz*, Amer. Math. Monthly **94** (1987), no. 4, 342–351.
- [62] Ranjan Roy, *Sources in the development of mathematics*, Cambridge University Press, 2011, Infinite series and products from the fifteenth to the twenty-first century.

- [63] Alexander A. Roytvarf, *Thinking in problems: How mathematicians find creative solutions*, Birkhäuser, 2013.
- [64] H. H. Schaefer, *Topological vector spaces*, second ed., Graduate Texts in Mathematics, vol. 3, Springer, 1999, With the assistance of M. P. Wolff.
- [65] Paul Schaefer, *Sums of rearranged series*, *College Math. J.* **17** (1986), no. 1, 66–70.
- [66] F. Smithies, *Cauchy's conception of rigour in analysis*, *Arch. Hist. Exact Sci.* **36** (1986), no. 1, 41–61.
- [67] Henrik Kragh Sørensen, *Throwing some light on the vast darkness that is analysis: Niels Henrik Abel's critical revision and the concept of absolute convergence*, *Centaurus* **52** (2010), no. 1, 38–72.
- [68] Lawrence E. Spence, *How many elements are in a union of sets?*, *The Mathematics Teacher* **80** (1987), no. 8, 666–670, 681.
- [69] John Tucciarone, *The development of the theory of summable divergent series from 1880 to 1925*, *Arch. Hist. Exact Sci.* **10** (1973), no. 1/2, 1–40.
- [70] Ian Tweddle, *James Stirling's early work on acceleration of convergence*, *Arch. Hist. Exact Sci.* **45** (1992), no. 2, 105–125.
- [71] ———, *James Stirling's methodus differentialis: An annotated translation of Stirling's text*, *Sources and Studies in the History of Mathematics and Physical Sciences*, Springer, 2003.
- [72] John Wallis, *The arithmetic of infinitesimals*, *Sources and Studies in the History of Mathematics and Physical Sciences*, Springer, 2004, Translated from the Latin and with an introduction by Jaequeline A. Stedall.
- [73] J. Watling, *The sum of an infinite series*, *Analysis* **13** (1952), no. 2, 39–46.
- [74] André Weil, *Elliptic functions according to Eisenstein and Kronecker*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 88, Springer, 1976.
- [75] Derek Thomas Whiteside, *Patterns of mathematical thought in the later seventeenth century*, *Arch. Hist. Exact Sci.* **1** (1961), no. 3, 179–388.
- [76] P. Wojtaszczyk, *A mathematical introduction to wavelets*, *London Mathematical Society Student Texts*, vol. 37, Cambridge University Press, 1997.
- [77] A. P. Youschkevitch, *The concept of function up to the middle of the 19th century*, *Arch. Hist. Exact Sci.* **16** (1976), no. 1, 37–85.